

Young functions on varifolds. Part I. Functional analytic foundations

Hsin-Chuang Chou

June 26, 2026

Abstract

The series of papers is devoted to the study of convergence for pairs of surfaces and smooth functions thereon. We model such pairs with varifolds and multiple-valued functions to capture their limits. In the present paper, we study Young functions, a measure-theoretic approach to multiple-valued functions, and the graph measures associated with pairs of measures (in particular, varifolds) and Young functions. This setting allows us to model the convergence of pairs of surfaces and functions thereon via the weak convergence of their associated graph measures, and a compactness theorem follows immediately. As a prerequisite for the concepts of differentiability for Young functions in the upcoming papers, we introduce and investigate several test function spaces.

Keywords: varifolds, multiple-valued functions, Young measures, Young functions, graph measures

MSC Classification: 28A35 (Primary), 46A13, 49Q20, 60B10 (Secondary)

Acknowledgements. The author would like to thank his PhD supervisor, Prof. Ulrich Menne, and his PhD co-supervisor, Dr. Nicolau Sarquis Aiex, for reading the paper and giving comments and suggestions. Also, the author would like to thank Mr. Yu-Tong Liu for consultation on functional analysis. The author was supported by the NTNU “Scholarship Pilot Program of the Ministry of Science and Technology to Subsidize Colleges and Universities in the Cultivation of Outstanding Doctoral Students”.

1 Introduction

This is the first of a series of papers aiming to answer the following questions.

Question. *In a Euclidean space, consider a sequence of smooth surfaces and smooth functions thereon such that over every compact subset the area of the surfaces and the integrals of mean curvature, the functions themselves, and their derivatives are uniformly bounded. What is the limit of these pairs? What can we say about the smoothness of the limit?*

Such questions typically arise when solving geometric variational problems, and those functions carry the geometric information about the surfaces. We

shall begin with the first question. Note that varifolds (see Allard [All72]) are a natural candidate to model the convergence of surfaces to retain information on their mean curvature in the limit process. With this model, we allow surfaces to have multiplicity, and the following example shows that we should also allow functions to have multiple values. Consider two parallel lines approaching each other, and on each of them, the functions have constant values 1 and -1 , respectively; in this case, the expected limit should consist of a line of multiplicity 2 and a constant two-valued function thereon. Almgren [Alm00] has developed (later extended by De Lellis and Spadaro [DLS11]) the theory of Q -valued functions in Euclidean spaces, where Q is a fixed positive integer, to study the regularity of minimizers of Plateau problems in higher codimensions. However, this theory is not quite sufficient for our setting because, in general, the behavior of varifold convergence is more complicated, the number of values of the limit function varies from one point to another, and the functions can oscillate such that the values of the limit function become diffuse. Therefore, we introduce *Young functions* (later or see 3.10) to model multiple-valued functions. Consequently, to answer the second question, we should develop the concepts of differentiability for Young functions on varifolds in advance.

The purpose of the present paper is to provide functional analytic foundations for later topics; more precisely, the concept of Young functions and the convergence of pairs of measures and Young functions. We also study some specific test function spaces on which the concepts of differentiability of Young functions rely. The upcoming papers will then contribute to, firstly, the study of differentiability for Young functions and the compatibility of our theory with that of Almgren's Q -valued functions; secondly, the compactness properties of pairs of varifolds and "differentiable" Young functions, and therefore the answer to the second question.

Description of the results. To present the results, we shall first introduce the concept of Young functions.

Definition (see 3.10). Suppose X and Y are locally compact Hausdorff spaces and μ is a Radon measure over X . By a μ *Young function* f of type Y , we mean a μ measurable $\mathbf{P}(Y)$ -valued function f , where $\mathbf{P}(Y)$ denotes the space of probability Radon measures over Y endowed with the initial topology induced from the maps $\nu \mapsto \int k d\nu$ for $\nu \in \mathbf{P}(Y)$ corresponding to continuous functions $k : Y \rightarrow \mathbf{R}$ with compact support.

Every μ measurable Y -valued function g gives rise to an associated μ Young function f of type Y such that $f(x) = \delta_{g(x)}$ for $x \in \text{dmn } g$, see 3.11; furthermore, we readily check that this statement still holds when g is a Lebesgue measurable Q -valued function by taking a measurable selection of g , see 3.13. Compared with these examples, we emphasize that the values of Young functions can be diffuse; also, the number of values can vary from one point to another. Less trivial examples of Young functions can be obtained from the Young measures generated by a sequence of functions in $\mathbf{L}_\infty(\mathcal{L}^m)$ and from the disintegration of measures on products (for instance, the disintegration of varifolds); the first example also justifies the terminology of Young functions, see 3.17 and 3.18.

To extend basic operations of single-valued functions to Young functions, we define typical operations on the space of probability measures, hence on the class of Young functions, such as pushforward, Cartesian product, and convolution, see 4.21, 4.22, and 4.27, respectively. When the Young functions are

associated with Y -valued functions, these operations then correspond to the post-composition, join¹, and addition (provided Y is a finite-dimensional Banach space) of Y -valued functions. Note that the notion of addition for multiple-valued functions is also new to the theory of Q -valued functions, but the addition of two Q -valued functions will be a Q^2 -valued function, see 4.24(3).

To study the convergence of pairs of measures and Young functions, we adapt Hutchinson's graph measures (see [Hut86, 4.3.1]) to our setting; in contrast to his concept of measure-function pairs in [Hut86, 4.1.1], we do not require any summability condition on Young functions.

Definition (see 3.14). Suppose X and Y are locally compact Hausdorff spaces, μ is a Radon measure over X , and f is a μ Young function of type Y . We define the *graph measure* $\mathbf{Y}(\mu, f)$ associated with μ and f to be the Radon measure over $X \times Y$ such that

$$\int \phi \, d\mathbf{Y}(\mu, f) = \int \phi(x, y) \, df(x) y \, d\mu x$$

whenever $\phi : X \times Y \rightarrow \mathbf{R}$ is a continuous function with compact support.

When X and Y have countable bases and f is the Young function associated with some μ measurable Y -valued function g , the graph measure $\mathbf{Y}(\mu, f)$ agrees with the pushforward of μ by the map $x \mapsto (x, g(x))$, see 3.16. The convergence of pairs is then defined to be the convergence of their associated graph measures. For the class of Radon measures Γ over $X \times Y$ such that $\Gamma(K \times Y) < \infty$ whenever K is a compact subset of X , we study its compactness property and present a disintegration theorem, see 3.23 and 3.25. Consequently, we have the following compactness theorem for pairs of varifolds and Young functions.

Theorem (see 3.27). *Suppose k, m , and n are positive integers, $m \leq n$, V_i is a sequence of m -dimensional varifolds in \mathbf{R}^n that converges to V , and f_i is a sequence of $\|V_i\|$ Young functions of type \mathbf{R}^k such that*

$$\lim_{s \rightarrow \infty} \sup \{ \mathbf{Y}(\|V_i\|, f_i)(K \times (\mathbf{R}^k \sim \mathbf{B}(0, s))) : i = 1, 2, \dots \} = 0$$

whenever K is a compact subset of \mathbf{R}^n . Then, there exists a $\|V\|$ Young function f of type \mathbf{R}^k such that, possibly passing to a subsequence, we have

$$\mathbf{Y}(\|V\|, f) = \lim_{i \rightarrow \infty} \mathbf{Y}(\|V_i\|, f_i).$$

Moreover, if we require V_i and V to be rectifiable, then we can further conclude that $\mathbf{Y}(V, f \circ p) = \lim_{i \rightarrow \infty} \mathbf{Y}(V_i, f_i \circ p)$, where $p : \mathbf{R}^n \times \mathbf{G}(n, m) \rightarrow \mathbf{R}^n$ is the projection map.

In preparation for investigating the concepts of differentiability for Young functions, we want to study the following test function spaces.

Definition (see 4.1). Suppose Y is a finite-dimensional Banach space. Denote by $\mathbf{E}_s(Y)$ the vector space of all continuously differentiable functions $\gamma : Y \rightarrow \mathbf{R}$ such that $\gamma(0) = 0$ and $\text{spt } D\gamma \subset \mathbf{B}(0, s)$. The space $\mathbf{E}_s(Y)$ is endowed with the norm whose value equals $\sup \text{im } \|D\gamma\|$ at $\gamma \in \mathbf{E}_s(Y)$, and we endow the vector space $\mathbf{E}(Y) = \bigcup_{0 \leq s < \infty} \mathbf{E}_s(Y)$ with the locally convex final topology (see 2.4) induced from the inclusions $\mathbf{E}_s(Y) \rightarrow \mathbf{E}(Y)$ corresponding to $0 \leq s < \infty$.

¹Suppose A, B , and C are sets and $f : A \rightarrow B$ and $g : A \rightarrow C$ are functions. By the join of f and g , we mean the function $A \rightarrow B \times C$ given by $a \mapsto (f(a), g(a))$.

Recall that a function on Euclidean space is termed weakly differentiable if and only if it satisfies an integration-by-parts identity. We may define weak differentiability of Y -valued functions g on varifolds in the same way, but as Menne pointed out in [Men16b, 8.27], the chain rule fails for the class of such functions. To resolve this problem, we have to require the compositions $\gamma \circ g$ for $\gamma \in \mathbf{E}(Y)$, instead of merely the functions g themselves, to satisfy an integration-by-parts identity, see [Men16b, 8.3]. When f is the Young function associated with g , we have $(\gamma \circ g)(x) = \int \gamma df(x)$ for $x \in \text{dmn } g$ and $\gamma \in \mathbf{E}(Y)$. Thus, the previous definition of weak differentiability suggests that it is expedient to view Young functions of type Y as $\mathbf{E}(Y)^*$ -valued functions, see 4.4. With this in mind, we study the topological vector space structure of $\mathbf{E}(Y)$ and give a homeomorphic embedding of $\mathbf{E}(Y)$ into a space of continuous functions with compact support, see 5.18. Moreover, to study the differentiability of Young functions on varifolds via their associated graph measures, we define the locally convex space $\mathbf{H}(U \times Y, \mathbf{R}^n)$ of certain test functions $\eta : U \times Y \rightarrow \mathbf{R}^n$, where U is an open subset of \mathbf{R}^n ; again, a homeomorphic embedding of $\mathbf{H}(U \times Y, \mathbf{R}^n)$ into a space of continuous functions with compact support is established, see 5.25.

As Lipschitz functions play an important role in the differentiation theory of functions on varifolds (cf. [Men16b] and [Men16a]), we expect their analogue for Young functions to play a similar role in our theory. To this end, we use $\mathbf{E}(Y)$ to define a pseudo-metric on $\mathbf{P}(Y)$ such that Dirac measures define an isometric embedding $Y \rightarrow \mathbf{P}(Y)$, which leads to a canonical definition of Lipschitz continuity for Young functions, see 4.7, 4.10, and 4.11. This pseudo-metric agrees with the 1-Wasserstein metric on the subspace $\mathbf{P}_1(Y)$ of those $\mu \in \mathbf{P}(Y)$ such that $\int |y| d\mu y < \infty$, and the metric topology on $\mathbf{P}_1(Y)$ is characterized by the weak convergence of measures μ and the convergence of their absolute first moments $\int |y| d\mu y$, see 4.14 and 4.15, respectively.

Organization of this paper. In Section 2, we collect the necessary materials for Section 5 from functional analysis. In Section 3, we define the notions of Young functions and their associated graph measures; also, we prove a compactness theorem for graph measures, a disintegration theorem, and therefore a compactness theorem for pairs of varifolds and Young functions. In Section 4, we study the space of probability Radon measures and define the operations push-forward, Cartesian product, and convolution on the class of Young functions. Moreover, we introduce Lipschitzian continuity for Young functions. In Section 5, we study the aforementioned test function spaces $\mathbf{E}(Y)$ and $\mathbf{H}(U \times Y, \mathbf{R}^n)$, for which embedding theorems into spaces of continuous functions with compact support are established.

Terminology. We employ the terms pseudo-metric, symmetric sets, absorbent sets, and locally convex spaces and locally convex topology, in accordance with Bourbaki, see [Bou89b, IX, §1, No. 1, Definition 1], [Bou89a, I, §4, No. 1, p. 31], [Bou87, I, §1, No. 5, Definition 4], [Bou87, II, §4, No. 1, Definition 1], respectively. We term a function of class k if and only if it is k -times continuously differentiable; moreover, we term a function of class ∞ if and only if it is of class k whenever k is a positive integer, see Federer [Fed69, 3.1.11].

Notation. The set of positive integers is denoted by \mathscr{P} . The set $\mathscr{P}^{\mathscr{P}}$ consists of all functions $\mathscr{P} \rightarrow \mathscr{P}$. If V is a vector space, $v \in V$, and $f \in \text{Hom}(V, \mathbf{R})$, then we denote $\langle v, f \rangle = f(v)$. If f is an $\overline{\mathbf{R}}$ -valued function defined on a directed set A , then its limit and limit superior (see Kelley [Kel75, p.

66]) are denoted by $\lim_{\alpha \in A} f(\alpha)$ and $\limsup_{\alpha \in A} f(\alpha)$, respectively. If F is the inductive limit of locally convex spaces F_α , see 2.2, then we denote $F = \varinjlim F_\alpha$. If U is an open subset of a finite-dimensional Banach space and Y is a Banach space, then $\mathcal{E}(U, Y)$ denotes the space of all functions of class ∞ from U into Y and $\mathcal{D}(U, Y) = \mathcal{E}(U, Y) \cap \{f : \text{spt } f \text{ is compact}\}$. The notation of Allard (see [All72]) will be employed for the discussion of varifolds.

Definitions in this paper. The notion of *directed sets* is defined in 2.1. The notions of *inductive systems* and their *inductive limits* are defined 2.2. The notions of *locally convex final topology* and *projective tensor product* are defined in 2.4 and 2.18, respectively. The space $\mathcal{K}(X, Z)$ of continuous functions with compact support and the space $\mathbf{P}(X)$ of probability Radon measures as well as their topologies are defined in 3.1 and 3.7, respectively. The notions of *Young functions* and their associated *graph measures* are defined in 3.10 and 3.14, respectively. The test function space $\mathbf{E}(Y)$ is introduced in 4.1. The pseudo-metric d on $\mathbf{P}(X)$ is defined in 4.7. The notion of *Lipschitzian Young functions* is defined in 4.11. The *pushforward* and *product* of Young functions are defined in 4.21 and 4.22, respectively. The *convolutions of measures* and the *convolution of Young functions* are defined in 4.23 and 4.27, respectively. The test function spaces $\tilde{\mathbf{E}}(Y)$ and $\mathbf{H}(U \times Y, \mathbf{R}^n)$ are introduced in 5.7 and 5.21, respectively.

2 Inductive limit and topological tensor product

In this section, we study the interaction between inductive limit and topological tensor product in the category of locally convex spaces.

Definition 2.1 (see [Kel75, p. 65]). We say a set A is *directed by the relation* \preceq if and only if \preceq is a reflexive and transitive relation on A such that every subset of A with at most two elements has an upper bound.

Definition 2.2 (see [Bou04b, III, §7, No. 5, p. 202] and [Bou87, II, §4, No. 4, Example II]). Suppose A is a set directed by the relation \preceq . An *inductive system of sets [resp. locally convex spaces] relative to A* consists of a family of sets [resp. locally convex spaces] F_α for $\alpha \in A$ and a family of maps [resp. continuous linear maps] $f_{\beta, \alpha}$ for $\alpha, \beta \in A$ with $\alpha \preceq \beta$ such that $f_{\alpha, \alpha} = \mathbf{1}_{F_\alpha}$ whenever $\alpha \in A$ and such that $f_{\gamma, \beta} \circ f_{\beta, \alpha} = f_{\gamma, \alpha}$ whenever $\alpha, \beta, \gamma \in A$ with $\alpha \preceq \beta \preceq \gamma$. In this case, the *inductive limit* of the inductive system $(F_\alpha, f_{\beta, \alpha})$ consists of a set [resp. a locally convex space] F and a family of maps [resp. continuous linear maps] f_α such that $f_\beta \circ f_{\beta, \alpha} = f_\alpha$ whenever $\alpha, \beta \in A$ with $\alpha \preceq \beta$, and satisfies the following universal property: whenever G is a set [resp. a locally convex space] and $g_\alpha : F_\alpha \rightarrow G$ is a map [resp. a continuous linear map] such that $g_\beta \circ f_{\beta, \alpha} = g_\alpha$ whenever $\alpha, \beta \in A$ with $\alpha \preceq \beta$, there exists a unique map [resp. a unique continuous linear map] $g : F \rightarrow G$ such that $g \circ f_\alpha = g_\alpha$ whenever $\alpha \in A$. For convenience, we will say F is the inductive limit of F_α for $\alpha \in A$ if either the maps $f_{\beta, \alpha}$ and f_α do not play a role in the statements or these maps are clear from the context.

Remark 2.3 (see [Bou04b, III, §7, No. 5, p. 202]). Suppose $(F_\alpha, f_{\beta, \alpha})$ forms an inductive system of sets relative to A . By convention, we identify F_α with its canonical image in the disjoint union $\bigsqcup_{\alpha \in A} F_\alpha$. Define F to be $\bigsqcup_{\alpha \in A} F_\alpha$ modulo the equivalence relation R such that $(x, y) \in R$ for some $x \in F_\alpha$ and $y \in F_\beta$ if

and only if there exists $\gamma \in A$ such that $\alpha \preceq \gamma, \beta \preceq \gamma$, and $f_{\gamma,\alpha}(x) = f_{\gamma,\beta}(y)$. For $\alpha \in A$, let f_α be the composition of the canonical maps $F_\alpha \rightarrow \bigsqcup_{\alpha \in A} F_\alpha \rightarrow F$. Then, the set F together with the maps f_α forms the inductive limit of the inductive system $(F_\alpha, f_{\beta,\alpha})$ of sets.

Definition 2.4 (see [Bou87, II, §4, No. 4, Proposition 5]). Suppose A is a set, F is a vector space, F_α is a locally convex space for $\alpha \in A$, and $f_\alpha : F_\alpha \rightarrow F$ is a linear map for $\alpha \in A$. Then, the *locally convex final topology* is the finest locally convex topology such that f_α is continuous whenever $\alpha \in A$. A fundamental system of neighborhoods of 0 of F is given by the family of all absorbent, convex, symmetric subsets V of F such that $f_\alpha^{-1}[V]$ is a neighborhood of 0 in F_α whenever $\alpha \in A$.

Remark 2.5 (see [Bou87, II, §4, No. 4, Proposition 5(ii)]). The locally convex final topology satisfies the following universal property: if G is a locally convex space and $L : F \rightarrow G$ is a linear map, then L is continuous if and only if $L \circ f_\alpha$ is continuous whenever $\alpha \in A$.

Remark 2.6 (see [Bou87, II, §4, No. 4, Example II]). Suppose A is a directed nonempty set, $(F_\alpha, f_{\beta,\alpha})$ forms an inductive system of locally convex spaces (hence an inductive system of sets) relative to A , and F and f_α are as in 2.3. Then, F can be endowed with a canonical vector structure such that each f_α is linear, see [Bou89a, II, §6, No. 2, p. 286]; moreover, if we endow F with the locally convex final topology induced by the maps f_α , then F together with f_α for $\alpha \in A$ forms the inductive limit of the inductive system $(F_\alpha, f_{\beta,\alpha})$ of locally convex spaces.

Remark 2.7 (see [Bou04b, III, §7, No. 6, Corollary 1 and Proposition 7]). Suppose A is a nonempty set directed by the relation \preceq , $(F_\alpha, f_{\beta,\alpha})$ and $(G_\alpha, g_{\beta,\alpha})$ are two inductive systems of sets [resp. locally convex spaces] relative to A , (F, f_α) and (G, g_α) are their inductive limits, respectively, and the maps [resp. continuous linear maps] $h_\alpha : F_\alpha \rightarrow G_\alpha$ for $\alpha \in A$ satisfy $h_\beta \circ f_{\beta,\alpha} = g_{\beta,\alpha} \circ h_\alpha$ whenever $\alpha, \beta \in A$ with $\alpha \preceq \beta$. By the universal property of inductive limits, there exists a unique map [resp. a unique continuous linear map] $h : F \rightarrow G$ such that $h \circ f_\alpha = g_\alpha \circ h_\alpha$ whenever $\alpha \in A$; in this case, h is called the inductive limit of h_α for $\alpha \in A$. Moreover, if each h_α is injective [resp. surjective], then h is injective [resp. surjective].

Definition 2.8 (see [Men16b, 2.9] and [Bou87, II, §4, No. 6, p. 33]). We say an inductive limit of locally convex spaces is *strict* if and only if the continuous linear maps in the corresponding inductive system are homeomorphic embeddings.

Remark 2.9. In the applications in Section 5, it will always be the case that F_α is a subspace of F_β whenever $\alpha \preceq \beta$, $F = \bigcup_{\alpha \in A} F_\alpha$, and the maps $f_{\beta,\alpha}$ and f_α are inclusion maps. In this case, F is the strict inductive limit of F_α if and only if F is endowed with the locally convex final topology induced by the inclusion maps f_α and each $f_{\beta,\alpha}$ is a homeomorphic embedding.

Lemma 2.10 (see [Bou87, II, §4, No. 6, Proposition 9] and [Bou87, III, §1, No. 4, Proposition 6]). Suppose $F = \bigcup_{i=1}^{\infty} F_i$ is the strict inductive limit of a sequence of Banach spaces F_i with $F_i \subset F_{i+1}$ for $i \in \mathcal{P}$. Then, the following two statements hold.

- (1) F is complete and Hausdorff, and the given topology on F_i agrees with the subspace topology induced by F ; in particular, F_i is a closed subspace of F .
- (2) Every bounded subset (in particular, compact subset) of F is contained in some F_i and bounded there.

Lemma 2.11 (see [Bou87, II, §4, No. 4, Corollary 2]). Suppose A and L are sets, $\{J_\lambda : \lambda \in L\}$ is a partition of A , G_α is a locally convex space for $\alpha \in A$, F_λ is a vector space for $\lambda \in L$, E is a vector space, $g_{\lambda,\alpha} : G_\alpha \rightarrow F_\lambda$ is a linear map for $\alpha \in J_\lambda$ and $\lambda \in L$, $h_\lambda : F_\lambda \rightarrow E$ is a linear map for $\lambda \in L$, and we endow F_λ with the locally convex final topology with respect to $g_{\lambda,\alpha}$ for $\alpha \in J_\lambda$ and $\lambda \in L$.

Then, the locally convex final topology on E induced by h_λ for $\lambda \in L$ agrees with the locally convex final topology induced by $h_\lambda \circ g_{\lambda,\alpha}$ for $\alpha \in J_\lambda$ and $\lambda \in L$.

Definition 2.12 (see [Bou87, II, §4, No. 5, Definition 2]). Suppose A is a set, F_α is a locally convex space for $\alpha \in A$, and F is the direct sum of F_α . Then, the direct sum topology on F is the locally convex final topology induced by the canonical injections $F_\alpha \rightarrow F$.

Lemma 2.13. Suppose A and B are sets, F_α^β and F_α for $(\alpha, \beta) \in A \times B$ are locally convex spaces, F_α is endowed with the locally convex final topology induced by the linear maps $f_\alpha^\beta : F_\alpha^\beta \rightarrow F_\alpha$ for $\beta \in B$ whenever $\alpha \in A$. Then, the locally convex final topology \mathcal{T} on $\bigoplus_{\alpha \in A} F_\alpha$ induced by the linear maps $\bigoplus_{\alpha \in A} f_\alpha^\beta$ for $\beta \in B$ agrees with the direct sum topology \mathcal{T}' of $\bigoplus_{\alpha \in A} F_\alpha$.

Proof. Denote the canonical injections by

$$i_\alpha : F_\alpha \rightarrow \bigoplus_{a \in A} F_a, \quad i_\alpha^\beta : F_\alpha^\beta \rightarrow \bigoplus_{a \in A} F_a^\beta.$$

In view of 2.11, \mathcal{T} is the locally convex final topology induced by the linear maps $(\bigoplus_{a \in A} f_a^\beta) \circ i_\alpha^\beta$ for $(\alpha, \beta) \in A \times B$, and \mathcal{T}' is the locally convex final topology induced by the linear maps $i_\alpha \circ f_\alpha^\beta$ for $(\alpha, \beta) \in A \times B$. On the other hand, we have

$$\left(\bigoplus_{a \in A} f_a^\beta \right) \circ i_\alpha^\beta = i_\alpha \circ f_\alpha^\beta$$

whenever $(\alpha, \beta) \in A \times B$. It follows that $\mathcal{T} = \mathcal{T}'$. \square

Lemma 2.14 (see [Bou87, II, §4, No. 3, Proposition 4]). Suppose A is a set, F_α is a locally convex space for $\alpha \in A$, F is a vector space, and $f_\alpha : F \rightarrow F_\alpha$ is a linear map for $\alpha \in A$. Then, the initial topology on F with respect to f_α for $\alpha \in A$ is a locally convex topology; in particular, the Cartesian product of locally convex spaces endowed with the product topology is again a locally convex space.

Remark 2.15. The following universal property of the Cartesian product of locally convex spaces follows from the corresponding property of topological spaces. Suppose A is a set, G and F_α for $\alpha \in A$ are locally convex spaces, F is the Cartesian product of F_α for $\alpha \in A$, f_α is the projection map for each $\alpha \in A$, and $g_\alpha : G \rightarrow F_\alpha$ is a continuous linear map for each $\alpha \in A$. Then, there exists a unique continuous linear map $h : G \rightarrow F$ such that $f_\alpha \circ h = g_\alpha$ whenever $\alpha \in A$.

Lemma 2.16 (see [Bou87, II, §4, No. 5, Proposition 7]). *Suppose A is a set, F_α is a locally convex space for $\alpha \in A$. Then, the canonical injection*

$$\bigoplus_{\alpha \in A} F_\alpha \rightarrow \prod_{\alpha \in A} F_\alpha$$

is continuous. If A is finite, then this map is an isomorphism of locally convex spaces.

Theorem 2.17. *Suppose (F, f_α) is the inductive limit [resp. the strict inductive limit] of the inductive system $(F_\alpha, f_{\beta, \alpha})$ of locally convex spaces. Then, $(F^n, \prod_{i=1}^n f_\alpha)$ is the inductive limit [resp. the strict inductive limit] of the inductive system $((F_\alpha)^n, \prod_{i=1}^n f_{\beta, \alpha})$ of locally convex spaces.*

Proof. By 2.14 and 2.15, we readily check that $((F_\alpha)^n, \prod_{i=1}^n f_{\beta, \alpha})$ form an inductive system of locally convex spaces. Observe that $(F^n, \prod_{i=1}^n f_\alpha)$ is the inductive limit of $((F_\alpha)^n, \prod_{i=1}^n f_{\beta, \alpha})$ in the category of sets. In view of 2.6, it is enough to show that F^n carries the locally convex final topology induced by the maps $\prod_{i=1}^n f_\alpha$ which is immediate by 2.16 and 2.13. To show the strictness of the inductive limit, we note that the Cartesian product of homeomorphic embeddings is again a homeomorphic embedding. \square

Definition 2.18 (see [SW99, III, 6.1 – 6.3]). Suppose V and W are locally convex spaces. The *projective tensor product topology* of $V \otimes W$ is the finest locally convex topology on $V \otimes W$ such that the natural bilinear map $\mu : V \times W \rightarrow V \otimes W$ is continuous.

Furthermore, if V and W are normed, then $V \otimes W$ is normed by the *projective tensor product norm* with value

$$\inf \left\{ \sum_{i=1}^n |v_i| |w_i| : \xi = \sum_{i=1}^n v_i \otimes w_i, v_i \in V, w_i \in W, i = 1, \dots, n \right\}$$

at $\xi \in V \otimes W$.

Remark 2.19. The projective tensor product topology satisfies the following property: whenever Z is a locally convex space and $g : V \times W \rightarrow Z$ is a continuous bilinear map, there exists a unique continuous linear map $h : V \otimes W \rightarrow Z$ such that $h = g \circ \mu$. In case that V, W and Z are normed, we have $\|g\| = \|h\|$.

Remark 2.20. Suppose $f_1 : V_1 \rightarrow W_1$ and $f_2 : V_2 \rightarrow W_2$ are continuous linear maps between locally convex spaces. Then, their tensor product $f_1 \otimes f_2 : V_1 \otimes V_2 \rightarrow W_1 \otimes W_2$ is continuous with respect to the projective tensor product topologies.

Remark 2.21. Suppose $V, W,$ and Z are normed spaces. It is straightforward to verify the following isometries of normed spaces

$$(V \otimes W) \otimes Z \simeq V \otimes (W \otimes Z), \quad V \otimes W \simeq W \otimes V.$$

If $\dim V < \infty$, we have the following isomorphism of normed spaces

$$\text{Hom}(V, \mathbf{R}) \otimes W \simeq \text{Hom}(V, W);$$

however, in general this is not an isometry.

Remark 2.22. Associating with a locally convex space V , the following isomorphism of locally convex spaces

$$I_V : \mathbf{R}^n \otimes V \simeq V^n.$$

yields a natural transformation; that is, whenever W is locally convex space and $f : V \rightarrow W$ is a continuous linear map, we have $I_W \circ (\mathbf{1}_{\mathbf{R}^n} \otimes f) = (\prod_{i=1}^n f) \circ I_V$.

Theorem 2.23. *Suppose (F, f_α) is the inductive limit [resp. the strict inductive limit] of the inductive system $(F_\alpha, f_{\beta, \alpha})$ of locally convex spaces. Then, $(\mathbf{R}^n \otimes F, \mathbf{1}_{\mathbf{R}^n} \otimes f_\alpha)$ is the inductive limit [resp. the strict inductive limit] of the inductive system $(\mathbf{R}^n \otimes F_\alpha, \mathbf{1}_{\mathbf{R}^n} \otimes f_{\beta, \alpha})$ of locally convex spaces.*

Proof. Combine 2.17 and 2.22. □

3 Young functions and graph measures

In this section, we introduce Young functions and graph measures, and we will focus on the compactness property of graph measures, leaving the discussion about Young functions to the next section.

We first collect necessary materials about the test function space $\mathcal{K}(X, Z)$ for later discussions on Young functions and graph measures.

Definition 3.1 (see [Bou04a, III, §1, No. 1]). Suppose X is a locally compact Hausdorff space and Z is a locally convex space. Then, the space $\mathcal{K}_K(X, Z)$ of continuous functions from X into Z with compact support in K is endowed with the supremum norm and the space

$$\mathcal{K}(X, Z) = \bigcup \{ \mathcal{K}_K(X, Z) : K \text{ is a compact subset of } X \}$$

is endowed with the locally convex final topology with respect to the inclusion maps $\mathcal{K}_K(X, Z) \rightarrow \mathcal{K}(X, Z)$ for compact subsets K of X . In case $Z = \mathbf{R}$, we will simply write $\mathcal{K}_K(X)$ and $\mathcal{K}(X)$; also, we denote $\mathcal{K}(X)^+ = \mathcal{K}(X) \cap \{f : f \geq 0\}$.

Remark 3.2. The topological dual space $\mathcal{K}(X)^*$ equals the space of all Daniell integrals on $\mathcal{K}(X)$, see [Fed69, 2.5.13], and it is endowed with the weak topology; that is, the initial topology induced from the maps $\mu \mapsto \mu(f)$ corresponding to $f \in \mathcal{K}(X)$, see [Fed69, 2.5.19]. For $\mu \in \mathcal{K}(X)^*$ such that $\mu(f) \geq 0$ whenever $f \in \mathcal{K}(X)^+$, there exists a unique Radon measure over X representing μ , see [Fed69, 2.5.13, 2.5.14].

Lemma 3.3. *Suppose X and Y are locally compact Hausdorff spaces, K and L are compact subsets of X and Y , respectively, Z is a Banach space. and the map*

$$\iota : Z^{X \times Y} \rightarrow (Z^Y)^X$$

is characterized by

$$\iota(\eta)(x) = \eta(x, \cdot) \quad \text{whenever } x \in X \text{ and } \eta \in Z^{X \times Y},$$

where B^A is the set of all functions $f : A \rightarrow B$. Then, the restriction

$$\iota|_{\mathcal{K}_{K \times L}(X \times Y, Z)} : \mathcal{K}_{K \times L}(X \times Y, Z) \rightarrow \mathcal{K}_K(X, \mathcal{K}_L(Y, Z))$$

defines a norm-preserving isomorphism of Banach spaces.

Proof. Clearly, we have $\sup \text{im} |\eta| = \sup \text{im} |\iota(\eta)|$ whenever $\eta \in \mathcal{K}_{K \times L}(X \times Y, Z)$. For topological spaces A and B , we denote by $\mathcal{C}(A, B)$ the space of all continuous functions $f : A \rightarrow B$ endowed with the compact-open topology. In case that B is a normed space, it is straightforward to verify that this topology agrees with the locally convex topology induced by the semi-norms with value $\sup |f|[K]$ at $f \in \mathcal{C}(A, B)$ corresponding to compact subsets K of A . It follows that the inclusion map $\mathcal{K}_K(X, B) \rightarrow \mathcal{C}(X, B)$ is a homeomorphic embedding whenever B is a normed space and K is a compact subset of X . Since the restriction

$$\iota|_{\mathcal{C}(X \times Y, Z)} : \mathcal{C}(X \times Y, Z) \rightarrow \mathcal{C}(X, \mathcal{C}(Y, Z))$$

defines a bijection by [tD08, 2.4.7], the assertion is obvious. \square

Remark 3.4. The above lemma is a generalization of [Bou04a, III, §4, No. 1, Lemma 1(i)] to vector-valued functions.

Lemma 3.5 (see [Bou04a, III, §1, No. 2, Proposition 5]). *Suppose X is a locally compact Hausdorff space, Z is a locally convex space, and K is a compact subset of X . Then, the image of the canonical map $\mathcal{K}_K(X) \otimes Z \rightarrow \mathcal{K}_K(X, Z)$ is dense.*

Lemma 3.6 (see [Bou04a, III, §4, No. 1, Lemma 1(ii)]). *Suppose X and Y are locally compact Hausdorff spaces, K and L are compact subsets of X and Y , respectively. Then, the image of the canonical map $\mathcal{K}_K(X) \otimes \mathcal{K}_L(Y) \rightarrow \mathcal{K}_{K \times L}(X \times Y)$ is dense.*

Proof. Apply 3.5 with $Z = \mathcal{K}_L(Y)$ and the assertion follows from 3.3. \square

Definition 3.7. Suppose X is a locally compact Hausdorff space. We define $\mathbf{P}(X)$ to be the space of all probability Radon measures over X , and it is endowed with the subspace topology induced from $\mathcal{K}(X)^*$.

Lemma 3.8. *Suppose X is a locally compact Hausdorff space. Then, the weak topology on $\mathbf{P}(X)$ agrees with the initial topology induced from the maps $\mu \mapsto \int g d\mu$ corresponding to bounded continuous functions $g : X \rightarrow \mathbf{R}$.*

Consequently, a sequence μ_i converges to μ weakly if and only if $\int g d\mu_i \rightarrow \int g d\mu$ whenever $g : X \rightarrow \mathbf{R}$ is a bounded continuous function.

Proof. Clearly, the weak topology of $\mathbf{P}(X)$ is weaker than the initial topology. Denoting by $e_g(\mu) = \int g d\mu$, then it is enough to show that $e_g : \mathbf{P}(X) \rightarrow \mathbf{R}$ is continuous with respect to the weak topology whenever $g : X \rightarrow \mathbf{R}$ is a non-negative bounded continuous function. By [Fed69, 2.2.5] and approximation with functions in $\mathcal{K}(X)$, we can show that

$$\mu(g) = \sup\{\mu(f) : 0 \leq f \leq g, f \in \mathcal{K}(X)\}$$

whenever $\mu \in \mathbf{P}(X)$ and $g : X \rightarrow \mathbf{R}$ is a non-negative bounded continuous function; in this case, it follows that e_g equals the supremum of the family of continuous functions e_f corresponding to $f \in \mathcal{K}(X)$ with $0 \leq f \leq g$, and therefore e_g is lower semi-continuous. Replacing g with $M - g$, where M is a large number such that $M - g \geq 0$, we have $e_g = -(e_{M-g} - M)$ is upper semi-continuous. \square

Remark 3.9. The initial topology mentioned in the lemma is usually termed weak topology in texts on probability theory (see [Kle20, 13.14(ii)]), and the lemma shows that these two notions of weak topology agree.

Definition 3.10. Suppose X and Y are locally compact Hausdorff spaces and μ is a Radon measure over X . By a μ Young function f of type Y , we mean a μ measurable function with values in $\mathbf{P}(Y)$.

Remark 3.11. Suppose $\delta : Y \rightarrow \mathcal{K}(Y)^*$ is given by $\delta_y(\beta) = \beta(y)$ whenever $y \in Y$ and $\beta \in \mathcal{K}(Y)$, g maps a subset of X into Y , $\mu(X \sim \text{dmn } g) = 0$, and $f = \delta \circ g$. Recall from [Fed69, 2.5.19] that δ is a homeomorphic embedding, it follows that g is μ measurable if and only if f is μ measurable.

Remark 3.12. In view of [Men16b, 2.12, 2.22], if Y is second-countable, then a $\mathbf{P}(Y)$ -valued function f defined for μ almost everywhere is μ measurable if and only if $x \mapsto \int \beta df(x)$ is μ measurable whenever $\beta \in \mathcal{K}(Y)$.

Example 3.13. Let $m, Q \in \mathcal{P}$. Suppose f is a \mathcal{L}^m measurable $\mathbf{Q}_Q(\mathbf{R}^n)$ -valued function in the sense of [Alm00, 1.1]. By [DLS11, 0.4], there exists \mathcal{L}^m measurable \mathbf{R}^n -valued functions f_1, f_2, \dots, f_Q such that $f = \delta \circ f_1 + \delta \circ f_2 + \dots + \delta \circ f_Q$; in particular, $Q^{-1}f$ is a $Q\mathcal{L}^m$ Young function of type \mathbf{R}^n by 3.12.

Definition 3.14. Suppose X and Y are locally compact Hausdorff spaces, μ is a Radon measure over X , and f is μ Young function of type Y . Then the graph measure $\mathbf{Y}(\mu, f)$ associated with μ and f is the Radon measure over $X \times Y$ such that (see 3.15)

$$\mathbf{Y}(\mu, f)(\psi) = \iint \psi(x, y) df(x) y d\mu x$$

whenever $\psi \in \mathcal{K}(X \times Y)$.

Remark 3.15. The notion of graph measure is well-defined. By 3.6, for $\psi \in \mathcal{K}(X \times Y)$ and $i \in \mathcal{P}$, there exist $k_i \in \mathcal{P}$, $\alpha_j \in \mathcal{K}(X)$ and $\beta_j \in \mathcal{K}(Y)$ for $j = 1, \dots, k_i$ such that

$$\left| \psi(x, y) - \sum_{j=1}^{k_i} \alpha_j(x) \beta_j(y) \right| \leq i^{-1} \quad \text{whenever } (x, y) \in X \times Y$$

and hence

$$\lim_{i \rightarrow \infty} \left| \int \psi(x, y) df(x) y - \sum_{j=1}^{k_i} \alpha_j(x) \int \beta_j(y) df(x) y \right| = 0 \quad \text{for } \mu \text{ almost all } x.$$

Thus, the map $x \mapsto \int \psi(x, y) df(x) y$ is μ measurable.

Remark 3.16. Suppose $m, n \in \mathcal{P}$ and g is a \mathcal{L}^m measurable function with values in \mathbf{R}^n . Then, $f = \delta \circ g$ is a \mathcal{L}^m Young function of type \mathbf{R}^n ; letting $G(x) = (x, g(x))$ for $x \in \text{dmn } g$, we have that $G_{\#}\mathcal{L}^m$ is a Radon measure by [MS25, 2.11] and [Fed69, 2.2.2], and that the graph measure $\mathbf{Y}(\mathcal{L}^m, f)$ associated with the pair (\mathcal{L}^m, f) equals $G_{\#}\mathcal{L}^m$. Compared with the notion of measure-function pairs introduced by Hutchinson (see [Hut86, 4.1.1]), our setting does not require any summability condition on functions g ; also, this example shows our notion of graph measure generalizes the one in [Hut86, 4.3.1].

Remark 3.17. Young functions are named after Laurence C. Young for their connection with Young measures. Suppose $m, n \in \mathcal{P}$ and U is a bounded open subset of \mathbf{R}^m . Using our terminology, the Young measures associated

with a bounded sequence of functions g_i in $\mathbf{L}_\infty(\mathcal{L}^m \llcorner U)$ form a $\mathcal{L}^m \llcorner U$ Young function f of type \mathbf{R}^n such that

$$\mathbf{Y}(\mathcal{L}^m \llcorner U, f) = \lim_{i \rightarrow \infty} \mathbf{Y}(\mathcal{L}^m \llcorner U, \delta \circ g_i),$$

see [AFP00, 2.30(i)] and 3.15.

Example 3.18. Suppose $m, n \in \mathcal{P}$ and U is an open subset of \mathbf{R}^n . Employing the notation of Allard's varifold disintegration (see [All72, 3.3]), for each varifold $V \in \mathbf{V}_m(U)$, the formula $x \mapsto V^{(x)}$ defines a $\|V\|$ Young function of type $\mathbf{G}(n, m)$ such that $V = \mathbf{Y}(\|V\|, V^{(\cdot)})$. A similar statement also holds for Young's generalized surfaces (which are termed oriented varifolds in [Hut86, Section 3]), see [Fle20, Part I, Section 4, (b), Section 10] and [Fle57, Section 3].

The following lemma shows that the disintegration formula in the definition of graph measures holds for a much larger class of functions.

Lemma 3.19. *Suppose X and Y are second-countable locally compact Hausdorff spaces, μ is a Radon measure over X , and f is a μ Young function of type Y . Then, we have*

$$\int \psi \, d\mathbf{Y}(\mu, f) = \iint \psi(x, y) \, df(x) y \, d\mu x$$

whenever ψ is a $\mathbf{Y}(\mu, f)$ integrable $\overline{\mathbf{R}}$ -valued function.

Proof. Let F consist of all $\mathbf{Y}(\mu, f)$ measurable subsets A of $X \times Y$ such that the formula remains true with ψ replaced by the characteristic function χ_A of A .

Step 1. Note that every open subset of $X \times Y$ is a countable union of compact subsets of $X \times Y$, and the disintegration formula holds for $\psi \in \mathcal{H}(X \times Y)$. In view of [Fed69, 2.4.7], F contains all the open subsets of $X \times Y$ and F is stable under countable increasing unions. If $A_i \in F$ for $i \in \mathcal{P}$ satisfies $\mathbf{Y}(\mu, f)(A_1) < \infty$ and $A_{i+1} \subset A_i$ for $i \in \mathcal{P}$, then $\bigcap_{i=1}^\infty A_i \in F$ by [Fed69, 2.4.9].

Step 2. If $\mathbf{Y}(\mu, f)(A) = 0$, there exists a countable nonempty subfamily $G \subset F$ of open subsets in $X \times Y$ such that $A \subset \bigcap G$ and $\mathbf{Y}(\mu, f)(\bigcap G) = 0$. Therefore, we have

$$\int^* \chi_A(x, \cdot) \, df(x) \leq \int \chi_{\bigcap G}(x, \cdot) \, df(x) = 0 \quad \text{for } \mu \text{ almost all } x$$

and this shows F contains all the sets A with $\mathbf{Y}(\mu, f)(A) = 0$.

Step 3. If A is a $\mathbf{Y}(\mu, f)$ measurable set with $\mathbf{Y}(\mu, f)(A) < \infty$, then there exists a countable nonempty subfamily $G \subset F$ of open subsets in $X \times Y$ such that $A \subset \bigcap G$ and $\mathbf{Y}(\mu, f)((\bigcap G) \sim A) = 0$, hence

$$\int \chi_{(\bigcap G) \sim A}(x, \cdot) \, df(x) = 0 \quad \text{for } \mu \text{ almost all } x;$$

it follows that

$$\int \chi_A(x, \cdot) \, df(x) = \int \chi_{\bigcap G}(x, \cdot) \, df(x) \quad \text{for } \mu \text{ almost all } x,$$

hence $A \in F$. Note that $X \times Y$ is countably $\mathbf{Y}(\mu, f)$ measurable. Therefore, F contains all $\mathbf{Y}(\mu, f)$ measurable sets.

Now, the assertion follows from [Fed69, 2.3.3, 2.4.8, 2.4.4(6)]. \square

To prove the compactness theorem and disintegration theorem, we want to reduce the problem by replacing Y with its one-point compactification Z ; for this purpose, we study the relation between the Radon measures over Y and their image under the inclusion map $\iota : Y \rightarrow Z$.

Lemma 3.20. *Suppose Z is a compact Hausdorff space, C is a closed subset of Z , and M is a nonempty compact family of Radon measures over Z such that $\phi(C) = 0$ for all $\phi \in M$. Then, there holds*

$$\limsup_{L \in P} \sup\{\phi(Z \sim L) : \phi \in M\} = 0,$$

where P is the family of compact subsets of $Z \sim C$ directed by the order \preceq such that $L_1 \preceq L_2$ if and only if $L_1 \subset L_2$ for $L_1, L_2 \in P$.

Proof. Let $\lambda = \limsup_{L \in P} \sup\{\phi(Z \sim L) : \phi \in M\}$ and let F consist of those $f \in \mathcal{X}(Z)$ satisfying $0 \leq f \leq 1$, and $C \subset \text{Int}\{z : f(z) = 1\}$. Since M is compact, whenever $f \in F$, there exists $\psi_f \in M$ such that

$$\psi_f(f) \geq \phi(f) \quad \text{whenever } \phi \in M;$$

furthermore, letting $L = Z \sim \text{Int}\{z : f(z) = 1\}$, we have $L \in P$ and $\psi_f(f) \geq \phi(Z \sim L)$ whenever $\phi \in M$, hence $\psi_f(f) \geq \lambda$. Observe that $\{\text{Clos}\{\psi_g : f \geq g \in F\} : f \in F\}$ forms a family of closed subsets of M that possesses the finite intersection property, then there exists

$$\phi \in \bigcap_{f \in F} \text{Clos}\{\psi_g : f \geq g \in F\}.$$

Since $\psi_g(f) \geq \psi_g(g) \geq \lambda$ and $f \mapsto \psi(f)$ for $\psi \in M$ is continuous whenever $f \geq g \in F$, we conclude $\phi(f) \geq \lambda$ whenever $f \in F$, thus $\lambda \leq \inf\{\phi(f) : f \in F\} = \phi(C) = 0$. \square

Lemma 3.21. *Suppose X and Y are locally compact Hausdorff spaces, Z is the one-point compactification of Y , and M is a family of Radon measures over $X \times Z$. Then, the following statements are equivalent.*

- (1) $\text{Clos } M$ is compact and $\phi(K \times (Z \sim Y)) = 0$ whenever K is a compact subset of X and $\phi \in \text{Clos } M$.
- (2) $\sup\{\phi(K \times Z) : \phi \in M\} < \infty$ whenever K is a compact subset of X and

$$\limsup_{L \in P} \sup\{\phi(K \times (Z \sim L)) : \phi \in M\} = 0,$$

where P is the family of compact subsets of Y directed by the order \preceq such that $L_1 \preceq L_2$ if and only if $L_1 \subset L_2$ for $L_1, L_2 \in P$.

Proof. To show (1) implies (2), we observe that for a compact subset K of X and a compact neighborhood G of K , there exists $f \in \mathcal{X}_{G \times Z}(X \times Z)^+$ such that $0 \leq f \leq 1$ and $K \times Z \subset \{x : f(x) = 1\}$, then mapping $\phi \in \text{Clos } M$ onto the finite Radon measure $(\phi \llcorner f)|_{\mathbf{2}^{G \times Z}}$ over $G \times Z$ defines a continuous map; from 3.20, we infer

$$\limsup_{L \in P} \sup\{\phi(K \times (Z \sim L)) : \phi \in M\} = 0.$$

Conversely, for a compact subset K of X , a compact neighborhood G of K , and $L \in P$, there exists $f_L \in \mathcal{K}(X \times Z)$ such that

$$0 \leq f_L \leq 1, \quad K \times (Z \sim Y) \subset \text{Int}\{x : f_L(x) = 1\}, \quad \text{spt } f_L \subset G \times (Z \sim L);$$

it follows that

$$\psi(K \times (Z \sim Y)) \leq \psi(f_L) \leq \sup\{\phi(G \times (Z \sim L)) : \phi \in M\},$$

hence $\psi(K \times (Z \sim Y)) = 0$ whenever $\psi \in \text{Clos } M$. \square

Lemma 3.22. *Suppose X and Y are locally compact Hausdorff spaces, Z is the one-point compactification of Y , $\iota : X \times Y \rightarrow X \times Z$ is the inclusion map, and the continuous monomorphism $j : \mathcal{K}(X \times Y) \rightarrow \mathcal{K}(X \times Z)$ is given by extension by zero to $X \times Z$. Then, there hold the following statements.*

- (1) *If X is a countable union of compact subsets and ψ is a Radon measure over $X \times Y$ such that $\psi(K \times Y) < \infty$ whenever K is a compact subset of X , then $\iota_{\#}\psi$ is a Radon measure over $X \times Z$.*
- (2) *If ϕ is a Radon measure over $X \times Z$ such that $\phi(K \times (Z \sim Y)) = 0$ whenever K is a compact subset of X and ψ is the Radon measure representing $j^*(\phi)$, then we have $\psi = \phi|_{\mathbf{2}^{X \times Y}}$ and ϕ is the Radon measure representing the member in $\mathcal{K}(X \times Z)^*$ induced by $\iota_{\#}\psi$.*
- (3) *Suppose M is a compact family of Radon measures ϕ over $X \times Z$ such that $\phi(K \times (Z \sim Y)) = 0$ whenever K is a compact subset of X . The dual map $j^* : \mathcal{K}(X \times Z)^* \rightarrow \mathcal{K}(X \times Y)^*$ embeds M homeomorphically into $\mathcal{K}(X \times Y)^*$.*

Proof. To prove (1), we first infer from [Fed69, 2.1.2] that every open subset of $X \times Z$ is $\iota_{\#}\psi$ measurable. Next, noting that $X \times Y$ is an open subset of $X \times Z$, we have

$$\begin{aligned} (\iota_{\#}\psi)(W) &= \psi(W \cap (X \times Y)) \\ &= \sup\{\psi(C) : C \text{ is compact, } C \subset W \cap (X \times Y)\} \\ &\leq \sup\{(\iota_{\#}\psi)(C) : C \text{ is compact, } C \subset W\} \end{aligned}$$

whenever W is an open subset of $X \times Z$. Let $\varepsilon > 0$ and let K_i be a sequence of compact subsets of X such that $X = \bigcup_{i=1}^{\infty} K_i$ and $K_i \subset \text{Int } K_{i+1}$ for $i \in \mathcal{P}$. Since ψ is a Radon measure and $\psi(K_i \times Y) < \infty$, there exists a sequence of compact subsets L_i of Y such that

$$\psi(K_i \times (Y \sim L_i)) < 2^{-i}\varepsilon.$$

Then $W = \bigcup_{i=1}^{\infty} (\text{Int } K_i) \times (Z \sim L_i)$ is an open superset of $X \times (Z \sim Y)$ such that $(\iota_{\#}\psi)(W) < \varepsilon$. For $A \subset X \times Z$, writing $A = (A \cap (X \times Y)) \cup (A \cap (X \times (Z \sim Y)))$, from the estimate above and the fact that ψ is a Radon measure, we conclude

$$(\iota_{\#}\psi)(A) = \inf\{(\iota_{\#}\psi)(W) : W \text{ is open, } A \subset W\}$$

whenever $A \subset X \times Z$.

As j is continuous, so is j^* . To prove (2), we compute by [Fed69, 2.4.18] that for $f \in \mathcal{K}(X \times Z)$ with $\text{spt } f \subset X \times Y$,

$$\int f d(\iota_{\#}\psi) = \int f \circ \iota d\psi = j^*(\phi)(f \circ \iota) = \phi(j(f \circ \iota)) = \int f d\phi.$$

Then, the first conclusion of (2) follows. For arbitrary $f \in \mathcal{K}(X \times Z)$, letting K be the image of projection of $\text{spt } f$ onto X , there exists a compact subset L of Y such that $\phi(K \times (Z \sim L))$ and $(\iota_{\#}\psi)(K \times (Z \sim L))$ are small; therefore, it allows us to approximate the integrals of f with the integrals of those functions whose supports are contained in $X \times Y$, and the second conclusion of (2) follows. By (2), $j^*|M$ is injective. Recalling that every continuous bijection from a compact space into a Hausdorff space is a homeomorphism, (3) follows. \square

Theorem 3.23 (Compactness). *Suppose X and Y are locally compact Hausdorff spaces, X is a countable union of compact subsets, and M is a class of Radon measures over $X \times Y$ satisfying*

$$\begin{aligned} \sup\{\Gamma(K \times Y) : \Gamma \in M\} &< \infty, \\ \limsup_{\Gamma \in P} \{\Gamma(K \times (Y \sim L)) : \Gamma \in M\} &= 0, \end{aligned}$$

whenever K is a compact subset of X , where P is the family of compact subsets of Y directed by the order \preceq such that $L_1 \preceq L_2$ if and only if $L_1 \subset L_2$ for $L_1, L_2 \in P$.

Then, $\text{Clos } M$ is compact, the push-forward $p_{\#}$ maps $\text{Clos } M$ continuously into $\mathcal{K}(X)^*$, and $p_{\#}\Gamma$ is a Radon measure over X whenever $\Gamma \in \text{Clos } M$, where $p : X \times Y \rightarrow X$ is the projection map.

Proof. Let Z be the one-point compactification of Y , let $\iota : X \times Y \rightarrow X \times Z$ be the inclusion map, let j be as in 3.22, and let $M' = \{\iota_{\#}\Gamma : \Gamma \in M\}$. From 3.22(1), the members of M' are Radon measures over $X \times Z$. Applying 3.21 with M replaced by M' , we see that $\text{Clos } M'$ is a compact family of Radon measures ϕ over $X \times Z$ satisfying $\phi(K \times (Z \sim Y)) = 0$ whenever K is a compact subset of X ; thus, $\text{Clos } M = j^*[\text{Clos } M']$ is compact and $\iota_{\#} \circ j^*| \text{Clos } M' = \mathbf{1}_{\text{Clos } M'}$ by 3.22(2)(3) applied with M replaced by $\text{Clos } M'$. By [Fed69, 2.2.17], $q_{\#}(\iota_{\#}\Gamma) = p_{\#}\Gamma$ is a Radon measure over X whenever $\Gamma \in \text{Clos } M$, where $q : X \times Z \rightarrow X$ is the projection map, and the continuity of $p_{\#}$ follows from the continuity of $(j^*| \text{Clos } M')^{-1}$. \square

Remark 3.24. The proof also shows that $\iota_{\#}$ maps $\text{Clos } M$ continuously into $\mathcal{K}(X \times Z)^*$. In particular, if $X \times Y$ is second-countable (and hence metrizable by [Kel75, Chapter 4, Theorem 16]) and M consists of a convergent sequence Γ_i and $\Gamma = \lim_{i \rightarrow \infty} \Gamma_i$, then we have $\iota_{\#}\Gamma = \lim_{i \rightarrow \infty} \iota_{\#}\Gamma_i$. It follows from [AFP00, 1.62] that

$$\lim_{i \rightarrow \infty} \int g d\Gamma_i = \int g d\Gamma$$

for any bounded Borel function g such that $\text{Clos } p[\text{spt } g]$ is compact and the set of discontinuity points of g has Γ measure zero.

Next, we shall present a disintegration theorem that suits our setting, see also [AFP00, 2.28].

Theorem 3.25. *Suppose X and Y are second-countable locally compact Hausdorff spaces, $p : X \times Y \rightarrow X$ is the projection map, and Γ is a Radon measure on $X \times Y$. If $\Gamma(K \times Y) < \infty$ for every compact subset K of X , then $p_{\#}\Gamma$ is a Radon measure over X and there exists a $p_{\#}\Gamma$ Young function f of type Y such that $\Gamma = \mathbf{Y}(p_{\#}\Gamma, f)$. Moreover, such a function f is $p_{\#}\Gamma$ almost unique.*

Proof. Suppose Z is the one-point compactification of Y , $\iota : Y \rightarrow Z$ is the inclusion map, and $q : X \times Z \rightarrow X$ is the projection map. Then Z is a second-countable compact Hausdorff space and by 3.22(1), $\mu = (\mathbf{1}_X \times \iota)_{\#}\Gamma$ is a Radon measure over $X \times Z$; in particular, $p_{\#}\Gamma = q_{\#}\mu$ is a Radon measure by [Fed69, 2.2.17]. Clearly, we have $\mu(X \times (Z \sim Y)) = 0$.

By 3.3, the function $S : \mathcal{K}(X, \mathcal{K}(Z)) \simeq \mathcal{K}(X \times Z)$ given by

$$S(u)(x, y) = u(x)(y) \quad \text{for } u \in \mathcal{K}(X, \mathcal{K}(Z))$$

is a linear isomorphism and apply [Fed69, 2.5.12] with $E = \mathcal{K}(Z)$ endowed with the sup-norm topology (which is separable by [Men16b, 2.2, 2.23]), $L = \mathcal{K}(X)$, $\Omega = \mathcal{K}(X, \mathcal{K}(Z))$, $T = \mu \circ S$, then there exists a Radon measure ϕ over X and a ϕ measurable function k with values in $\mathcal{K}(Z)^*$ with respect to the weak topology such that $|k(x)| = 1$ for ϕ almost all x and such that

$$\int S(u) d\mu = T(u) = \int \langle u(x), k(x) \rangle d\phi x \quad \text{whenever } u \in \mathcal{K}(X, \mathcal{K}(Z)).$$

For $\beta \in \mathcal{K}(Z)^+$, we have

$$0 \leq \int \alpha(x)\beta(z) d\mu(x, z) = \int \alpha(x)\langle \beta, k(x) \rangle d\phi x \quad \text{whenever } \alpha \in \mathcal{K}(X)^+;$$

thus, $\langle \beta, k(x) \rangle \geq 0$ for ϕ almost all x . As $\mathcal{K}(Z)^+$ is separable, we conclude for ϕ almost all x , $k(x)$ is monotone, hence $1 = |k(x)| = k(x)(Z)$. Therefore, k is a ϕ Young function of type Z and $\mu = \mathbf{Y}(\phi, k)$, hence $p_{\#}\Gamma = q_{\#}\mu = \phi$. By 3.19, we conclude that $k(x)(Z \sim Y) = 0$ for ϕ almost all x .

Note that extension by zero gives a continuous monomorphism

$$j : \mathcal{K}(Y) \rightarrow \mathcal{K}(Z),$$

hence a continuous linear map

$$j^* : \mathcal{K}(Z)^* \rightarrow \mathcal{K}(Y)^*,$$

where $\mathcal{K}(Z)^*$ and $\mathcal{K}(Y)^*$ are endowed with the weak topology. Therefore, $f = j^* \circ k$ is a $p_{\#}\Gamma$ Young function of type Y such that

$$\begin{aligned} \int \psi d\Gamma &= \int j(\psi(x, \cdot))(y) d\mu(x, y) \\ &= \int \langle j(\psi(x, \cdot)), k(x) \rangle d\phi x \\ &= \int \langle \psi(x, \cdot), f(x) \rangle d\phi x \\ &= \iint \psi(x, y) df(x) y d(p_{\#}\Gamma) x \end{aligned}$$

whenever $\psi \in \mathcal{K}(X \times Y)$.

Finally, suppose g is a $p_{\#}\Gamma$ Young function of type Y such that

$$\mathbf{Y}(p_{\#}\Gamma, g) = \mathbf{Y}(p_{\#}\Gamma, f).$$

By 3.8 and 3.22(1), we see $x \mapsto \iota_{\#}(g(x))$ for $x \in \text{dmn } g$ is a $p_{\#}\Gamma$ Young function of type Z . Note that $k(x)(Z \sim Y) = 0$, hence by 3.22(2), $\iota_{\#}(f(x)) = k(x)$ for $p_{\#}\Gamma$ almost all x . By 3.19 and [Fed69, 2.4.18], we compute for $u \in \mathcal{K}(X, \mathcal{K}(Z))$

$$\begin{aligned}
T(u) &= \int \langle u(x), k(x) \rangle d(p_{\#}\Gamma) x \\
&= \iint S(u)(x, \iota(y)) df(x) y d(p_{\#}\Gamma) x \\
&= \int S(u)(x, \iota(y)) d\mathbf{Y}(p_{\#}\Gamma, f)(x, y) \\
&= \int S(u)(x, \iota(y)) d\mathbf{Y}(p_{\#}\Gamma, g)(x, y) \\
&= \iint S(u)(x, \iota(y)) dg(x) y d(p_{\#}\Gamma) x \\
&= \iint S(u)(x, z) d\iota_{\#}(g(x)) z d(p_{\#}\Gamma) x \\
&= \int \langle u(x), \iota_{\#}(g(x)) \rangle d(p_{\#}\Gamma) x.
\end{aligned}$$

From the uniqueness of k , we conclude $\iota_{\#}(g(x)) = k(x)$, hence by 3.22(2), $g(x) = j^*(\iota_{\#}(g(x))) = (j^* \circ k)(x) = f(x)$ for $p_{\#}\Gamma$ almost all x . \square

Finally, we will finish this section with the compactness theorem for pairs of varifolds and Young functions.

The following lemma shows that if V is a rectifiable varifold, then there is a one-to-one correspondence between $\|V\|$ Young functions and V Young functions obtained from precomposition with the functions τ and p as in the lemma.

Lemma 3.26. *Suppose m and n are positive integers, $m \leq n$, V is an m -dimensional varifold in some open set U of \mathbf{R}^n , and define τ by*

$$\tau(x) = (x, \text{Tan}^m(\|V\|, x)) \quad \text{whenever } \text{Tan}^m(\|V\|, x) \in \mathbf{G}(n, m).$$

Then, the following statements hold.

- (1) $V = \tau_{\#}\|V\|$.
- (2) For $A \subset U \times \mathbf{G}(n, m)$, A is V measurable if and only if $\tau^{-1}[A]$ is $\|V\|$ measurable.
- (3) $\tau(p(x, S)) = (x, S)$ for V almost all (x, S) , where $p : U \times \mathbf{G}(n, m) \rightarrow U$ is the projection map.

Proof. By [MS25, 2.11] and [All72, 3.5(1b)], $\tau_{\#}\|V\|$ is a Radon measure and $V = \tau_{\#}\|V\|$. Let $A \subset U \times \mathbf{G}(n, m)$. Clearly, if $\tau^{-1}[A]$ is $\|V\|$ measurable, then A is $V = \tau_{\#}\|V\|$ measurable. If A is V measurable, then there exists a Borel set B such that $A \subset B$ and $V(B \sim A) = 0$, hence $\|V\|\tau^{-1}[B \sim A] = 0$. Therefore,

$$\tau^{-1}[A] = \tau^{-1}[B] \sim \tau^{-1}[B \sim A]$$

is $\|V\|$ measurable and (2) follows. To prove (3), we note that

$$\tau^{-1}[\{(x, S) : (x, S) \neq (x, \text{Tan}^m(\|V\|, x))\}] = \emptyset,$$

and the assertion follows from (1). \square

Theorem 3.27 (Compactness). *Suppose m and n are positive integers, $m \leq n$, V_i is a sequence of m -dimensional varifolds in some open set U of \mathbf{R}^n such that $V = \lim_{i \rightarrow \infty} V_i$, Y is a second-countable locally compact Hausdorff space, L_j is*

a sequence of compact subsets of Y such that $\bigcup_{j=1}^{\infty} L_j = Y$ and $L_j \subset \text{Int } L_{j+1}$ whenever $j \in \mathcal{P}$, and f_i is a sequence of $\|V_i\|$ Young functions of type Y such that

$$\lim_{j \rightarrow \infty} \sup \{ \mathbf{Y}(\|V_i\|, f_i)(K \times (Y \sim L_j)) : i \in \mathcal{P} \} = 0,$$

whenever K is a compact subset of U .

Then, exists a $\|V\|$ Young function f of type Y such that, possibly passing to a subsequence,

$$\mathbf{Y}(\|V\|, f) = \lim_{i \rightarrow \infty} \mathbf{Y}(\|V_i\|, f_i);$$

moreover, if V_i and V are rectifiable, then

$$\mathbf{Y}(V, f \circ p) = \lim_{i \rightarrow \infty} \mathbf{Y}(V_i, f_i \circ p),$$

where $p : U \times \mathbf{G}(n, m) \rightarrow U$ is the projection map.

Proof. The main assertion follows from 3.23 and 3.25, and the postscript follows from 3.26. \square

4 The space of probability Radon measures and operations on Young functions

In this section, we first define a metric on the space of probability Radon measures and study its topological and metric structures. Then, we show that the class of Young functions is stable under several operations.

Definition 4.1. Suppose Y is a finite-dimensional Banach space. For $0 \leq s < \infty$, the space $\mathbf{E}_s(Y)$ consists of functions $\gamma : Y \rightarrow \mathbf{R}$ of class 1 such that $\gamma(0) = 0$ and $\text{spt } D\gamma \subset \mathbf{B}(0, s)$. We endow $\mathbf{E}_s(Y)$ with the norm whose value equals

$$\sup \text{im } \|D\gamma\| \quad \text{at } \gamma \in \mathbf{E}_s(Y),$$

and we endow $\mathbf{E}(Y) = \bigcup \{ \mathbf{E}_s(Y) : 0 \leq s < \infty \}$ with the locally convex final topology induced by the inclusion maps $\mathbf{E}_s(Y) \rightarrow \mathbf{E}(Y)$.

Remark 4.2. Suppose Y is a finite-dimensional Banach space. Letting $k = \dim Y$ and employing the linear isomorphism $Y \simeq \mathbf{R}^k$, the standard mollification argument in Euclidean spaces also works for Y .

Remark 4.3. Whenever $0 \leq s < \infty$ and $\gamma \in \mathbf{E}_s(Y)$, we have

$$|\gamma(y)| \leq \inf\{|y|, s\} \sup \text{im } \|D\gamma\| \quad \text{for } y \in Y;$$

in particular, γ is bounded.

Remark 4.4. The canonical map $\mathbf{P}(Y) \rightarrow \mathbf{E}(Y)^*$ is continuous by 3.8 and 4.3; so is its composition with the dual map $\mathbf{E}(Y)^* \rightarrow \mathbf{E}_s(Y)^*$ whenever $0 \leq s < \infty$, where the dual spaces are endowed with the corresponding weak topology.

The following lemma shows that, with the help of Ascoli's theorem, the condition on the support of functions in $\mathbf{E}_s(Y)$ allows us to retain the continuity of the canonical map $\mathbf{P}(Y) \rightarrow \mathbf{E}_s(Y)^*$ with respect to the dual norm topology on $\mathbf{E}_s(Y)^*$.

Lemma 4.5. *Suppose Y is a finite-dimensional Banach space and $0 \leq s < \infty$. Then, the canonical map $\mathbf{P}(Y) \rightarrow \mathbf{E}_s(Y)^*$ is continuous with bounded image, where $\mathbf{E}_s(Y)^*$ is endowed with the dual norm topology.*

Proof. By [Fed69, 2.10.21], the function space

$$\mathbf{E}_s(Y) \cap \{\gamma : \sup \text{im} \|\mathbf{D}\gamma\| \leq 1\}$$

is contained in the compact space F of Lipschitzian functions $f : Y \rightarrow \mathbf{R}$ with $f(0) = 0$ and $\text{Lip } f \leq 1$, where F is endowed with the topology of uniform convergence on each compact subset of Y , or equivalently, the topology induced by the metric ρ with value

$$\rho(f, g) = \sum_{i=1}^{\infty} 2^{-i} \cdot \frac{\sup \text{im} |(f - g)|_{\mathbf{B}(0, i)}}{1 + \sup \text{im} |(f - g)|_{\mathbf{B}(0, i)}} \quad \text{at } (f, g) \in F \times F.$$

Note that

$$\sup \text{im} |\gamma_1 - \gamma_2| = \sup\{|\gamma_1(y) - \gamma_2(y)| : y \in Y \cap \mathbf{B}(0, s)\}$$

whenever $\gamma_1, \gamma_2 \in \mathbf{E}_s(Y)$. Then, it is straightforward to show that the function space $\mathbf{E}_s(Y) \cap \{\gamma : \sup \text{im} \|\mathbf{D}\gamma\| \leq 1\}$ is totally bounded with respect to the supremum metric. With the help of 3.8, it follows that every convergent sequence in $\mathbf{P}(Y)$ also converges in $\mathbf{E}_s(Y)^*$. Since $\mathbf{P}(Y) \subset \mathcal{K}(Y)^*$ is metrizable by [Men16b, 2.23], the continuity of the canonical map follows, and the boundedness of its image is immediate by 4.3. \square

Lemma 4.6. *Suppose Y is a finite-dimensional Banach space. There holds*

$$|y_1 - y_2| = \sup\{\gamma(y_1) - \gamma(y_2) : \gamma \in \mathbf{E}(Y), \sup \text{im} \|\mathbf{D}\gamma\| \leq 1\}$$

whenever $y_1, y_2 \in Y$.

Proof. Since $|\gamma(y_1) - \gamma(y_2)| \leq |y_1 - y_2| \sup \text{im} \|\mathbf{D}\gamma\|$ whenever $\gamma \in \mathbf{E}(Y)$, it suffices to show there exists a sequence $\gamma_i \in \mathbf{E}(Y)$ such that

$$\lim_{i \rightarrow \infty} \gamma_i(y_1) - \gamma_i(y_2) = |y_1 - y_2|.$$

Note that there exists $L \in \text{Hom}(Y, \mathbf{R})$ such that $L(y_1 - y_2) = |y_1 - y_2|$ and $\|L\| \leq 1$. Whenever $i \in \mathcal{P}$, since $L_i = \sup\{\inf\{L, i\}, -i\}$ is uniformly continuous and $L_i(0) = 0$, there exists $\beta_i : Y \rightarrow \mathbf{R}$ of class 1 such that $\beta_i(0) = 0$, $\sup \text{im} \|\mathbf{D}\beta_i\| \leq 1$, and $\sup \text{im} |L_i - \beta_i| \leq i^{-1}$. Choose a function $\phi : Y \rightarrow \mathbf{R}$ of class 1 such that $0 \leq \phi \leq 1$, $\phi|_{\mathbf{B}(0, 1)} = 1$, $\sup \text{im} \|\mathbf{D}\phi\| \leq 1$, and $\text{spt } \phi$ is compact, and we define $\phi_i(y) = (1 - i^{-1})\phi(i^{-3}y)$ whenever $y \in Y$ and $i \in \mathcal{P}$. Then, we have $\gamma_i = \phi_i\beta_i \in \mathbf{E}(Y)$, $\sup \text{im} \|\mathbf{D}\gamma_i\| \leq 1$ for $i \geq 1$, and $L(y) = \lim_{i \rightarrow \infty} \gamma_i(y)$ for all $y \in Y$ and the assertion follows. \square

Definition 4.7. Suppose Y is a finite-dimensional Banach space, we define the pseudo-metric d on $\mathbf{P}(Y)$ by

$$d(\mu, \nu) = \sup \left\{ \int \gamma \, d\mu - \int \gamma \, d\nu : \gamma \in \mathbf{E}(Y), \sup \text{im} \|\mathbf{D}\gamma\| \leq 1 \right\}$$

whenever $\mu, \nu \in \mathbf{P}(Y)$.

Remark 4.8. If $\mu, \nu \in \mathbf{P}(Y)$ satisfy $d(\mu, \nu) = 0$, we have $\int \gamma d\mu = \int \gamma d\nu$ whenever $\gamma \in \mathcal{K}(Y)$ with $\text{Lip } \gamma < \infty$; it follows that $\mu = \nu$.

Remark 4.9. By approximation, we readily show that

$$\begin{aligned} d(\mu, \nu) &= \sup \left\{ \int \gamma d\mu - \int \gamma d\nu : \gamma : Y \rightarrow \mathbf{R}, \text{Lip } \gamma \leq 1 \right\} \\ &= \sup \left\{ \int \gamma d\mu - \int \gamma d\nu : \gamma \in \mathcal{K}(Y), \text{Lip } \gamma \leq 1 \right\}. \end{aligned}$$

whenever $\mu, \nu \in \mathbf{P}(Y)$.

Remark 4.10. It follows from 4.6 that $\delta : Y \rightarrow \mathbf{P}(Y)$ as in 3.11 is an isometric embedding.

Definition 4.11. Suppose (X, ρ) is a pseudo-metric space and Y is a finite-dimensional Banach space. A function $f : X \rightarrow \mathbf{P}(Y)$ is termed to be *Lipschitzian* if and only if there exists $0 < M < \infty$ such that

$$d(f(x), f(y)) \leq M\rho(x, y) \quad \text{whenever } x, y \in X,$$

where d as in 4.7. The Lipschitz constant of f is defined to be the infimum of all such numbers M . We say f is *locally Lipschitzian* if and only if whenever $x \in X$, there exists a neighborhood U of x such that $f|_U$ is Lipschitzian.

Remark 4.12. Note that if X is a metric space and f is a Lipschitzian $\mathbf{P}(Y)$ -valued function, then we have $d(f(x), f(y)) < \infty$ whenever $x, y \in X$; in particular, $d|(\text{im } f \times \text{im } f)$ is a metric and f is Lipschitzian in the usual sense.

Remark 4.13. By 4.9, we have

$$d(\delta_0, \mu) = \int |y| d\mu y \quad \text{whenever } \mu \in \mathbf{P}(Y).$$

Let

$$\mathbf{P}_1(Y) = \mathbf{P}(Y) \cap \left\{ \mu : \int |y| d\mu y < \infty \right\}.$$

Then, $(\mathbf{P}_1(Y), d)$ is a metric space.

Next, we shall investigate the relation between the weak topology and the d topology on $\mathbf{P}_1(Y)$ when Y is a finite-dimensional Banach space.

Remark 4.14. The notation $\mathbf{P}_1(Y)$ is taken from [Vil09, 6.4], on which the metric d agree with 1-Wasserstein metric, see [Vil09, 6.1, 6.5]. The following characterization of d convergence on $\mathbf{P}_1(Y)$ is a special case of [Vil09, 6.9].

Lemma 4.15 (see [Vil09, 6.8, 6.9]). *Suppose Y is a finite-dimensional Banach space, μ_i is a sequence in $\mathbf{P}_1(Y)$, and $\mu \in \mathbf{P}_1(Y)$. Then, the following two statements are equivalent.*

- (1) $\lim_{i \rightarrow \infty} d(\mu_i, \mu) = 0$.
- (2) $\mu_i \rightarrow \mu$ weakly as $i \rightarrow \infty$ and $\lim_{i \rightarrow \infty} \int |y| d\mu_i y = \int |y| d\mu y$.

Proof. Suppose $\lim_{i \rightarrow \infty} d(\mu_i, \mu) = 0$, then

$$\lim_{i \rightarrow \infty} \int |y| d\mu_i y = \lim_{i \rightarrow \infty} d(\delta_0, \mu_i) = d(\delta_0, \mu) = \int |y| d\mu y.$$

Whenever $\varepsilon > 0$ and $f \in \mathcal{K}(Y)$, since f is uniformly continuous, there exists a function $\gamma : Y \rightarrow \mathbf{R}$ of class 1 such that $\text{spt } \gamma$ is compact and $\sup \text{im } |f - \gamma| < \varepsilon$, hence we have

$$|\mu_i(f) - \mu(f)| \leq 2\varepsilon + |\mu_i(\gamma) - \mu(\gamma)| \leq 2\varepsilon + \sup \text{im } \|D\gamma\| d(\mu_i, \mu).$$

Therefore, we conclude $\mu_i \rightarrow \mu$ weakly as $i \rightarrow \infty$.

To prove the converse, letting $\varepsilon > 0$, we will first show that there exists $0 \leq s < \infty$ such that $\int_{Y \sim \mathbf{B}(0,s)} |y| d\mu y \leq \varepsilon$ and

$$\int_{Y \sim \mathbf{B}(0,s)} |y| d\mu_i y \leq \varepsilon \quad \text{whenever } i \in \mathcal{P}.$$

Since μ is a Radon measure, there exists $f \in \mathcal{K}(Y)$ such that $0 \leq f \leq 1$ and $\int (1 - f(y)) |y| d\mu y \leq \varepsilon/2$. From the hypothesis, there exists $N \in \mathcal{P}$ such that

$$|\int f(y) |y| d\mu_i y - \int f(y) |y| d\mu y| + |\int |y| d\mu_i y - \int |y| d\mu y| \leq \varepsilon/2$$

whenever $i \geq N$. Then, we have

$$\int_{Y \sim \text{spt } f} |y| d\mu_i y \leq \varepsilon \quad \text{whenever } i \geq N$$

and the assertion follows. On the other hand, by [Fed69, 2.10.21], the set $\mathcal{K}_{\mathbf{B}(0,r)}(Y) \cap \{\gamma : \text{Lip } \gamma \leq 1\}$ is compact in the Banach space $\mathcal{K}_{\mathbf{B}(0,r)}(Y)$; since each member of $\mathbf{P}(Y)$ defines a continuous linear functional over $\mathcal{K}_{\mathbf{B}(0,r)}$, we have

$$\limsup_{i \rightarrow \infty} \{\int \gamma d\mu_i - \int \gamma d\mu : \gamma \in \mathcal{K}_{\mathbf{B}(0,r)}(Y), \text{Lip } \gamma \leq 1\} = 0$$

whenever $0 < r < \infty$. It follows that $\lim_{i \rightarrow \infty} d(\mu_i, \mu) = 0$. \square

Theorem 4.16. *The map $(\mathbf{P}_1(Y), d) \rightarrow \mathbf{P}(Y) \times \mathbf{R}$ defined by*

$$\mu \mapsto (\mu, \int |y| d\mu y) \quad \text{for } \mu \in \mathbf{P}_1(Y)$$

is a homeomorphic embedding.

Proof. Since the weak topology on the space of Radon measures over Y is metrizable by [Men16b, 2.23], the assertion follows from 4.15. \square

Remark 4.17. We can not replace $\mathbf{P}_1(Y)$ with $\mathbf{P}(Y)$ in 4.15. To construct a counterexample, taking $Y = \mathbf{R}$, $\mu \in \mathbf{P}(\mathbf{R}) \sim \mathbf{P}_1(\mathbf{R})$, and $\mu_i = \mu \llcorner \mathbf{B}(0, i)$, it is easy to verify $\mu_i \rightarrow \mu$ weakly and $\int |y| d\mu_i y \uparrow \int |y| d\mu y$ as $i \rightarrow \infty$. Since $\mu_i \in \mathbf{P}_1(\mathbf{R})$ and $\mu \in \mathbf{P}(\mathbf{R}) \sim \mathbf{P}_1(\mathbf{R})$, we have $d(\mu_i, \delta_0) < \infty$ and $d(\mu, \delta_0) = \infty$ by 4.13; it follows that $d(\mu_i, \mu) = \infty$ for all $i \in \mathcal{P}$.

Remark 4.18. The d topology is strictly finer than the weak topology on $\mathbf{P}_1(Y)$ provided $\dim Y \geq 1$; for instance, letting $0 \neq v \in Y$ and $\mu \in \mathbf{P}_1(Y)$, the sequence $\mu_i = (1 - i^{-1})\mu + i^{-1}\delta_{iv}$ converges weakly to μ as $i \rightarrow \infty$ but

$$\lim_{i \rightarrow \infty} \int |y| d\mu_i y - \int |y| d\mu y = |v| > 0.$$

However, these two topologies induce the same subspace topology on the weakly closed subsets

$$A(\varepsilon) = \mathbf{P}(Y) \cap \{\mu : \mu\{a\} \geq \varepsilon \text{ whenever } a \in \text{spt } \mu\} \subset \mathbf{P}_1(Y)$$

whenever $\varepsilon > 0$. To show this, suppose $\varepsilon > 0$ and μ_i is a sequence in $A(\varepsilon)$ that converges weakly to some $\mu \in \mathbf{P}(Y)$. Whenever $0 < r < \infty$ and $a \in \text{spt } \mu$, by [All72, 2.6(2c)], there exists $N \in \mathcal{P}$ such that $\mu_i(\mathbf{U}(a, r)) > 0$, hence $\mu_i(\mathbf{U}(a, r)) \geq \varepsilon$, whenever $i \geq N$; it follows that

$$\mu(\mathbf{B}(a, r)) \geq \limsup_{i \rightarrow \infty} \mu_i(\mathbf{B}(a, r)) \geq \varepsilon.$$

Thus, we conclude $\mu \in A(\varepsilon)$. Since $\lim_{i \rightarrow \infty} \int (1-f) d\mu_i = \int (1-f) d\mu$ whenever $f \in \mathcal{K}(Y)$ and note that $\nu(S) > 0$ implies $\nu(S) \geq \varepsilon$ whenever $S \subset Y$ and $\nu \in A(\varepsilon)$, there exists $0 < s < \infty$ such that

$$\sup\{\mu_i(Y \sim \mathbf{B}(0, s)) : i \in \mathcal{P}\} = 0 \quad \text{for some } 0 < s < \infty;$$

by 3.8,

$$\lim_{i \rightarrow \infty} \int |y| d\mu_i y = \lim_{i \rightarrow \infty} \int \inf\{|y|, s\} d\mu_i y = \int \inf\{|y|, s\} d\mu y = \int |y| d\mu y.$$

The following theorem shows that the class of Young functions is stable under pushforward by Borel functions pointwise and taking the Cartesian product of measures pointwise.

Theorem 4.19. *Suppose X and Y are second-countable locally compact Hausdorff spaces and $f : X \rightarrow Y$ is a Borel function. Then, the following two statements hold.*

(1) *The pushforward of measures*

$$(f_{\#}\mu)(A) = \mu(f^{-1}[A]) \quad \text{whenever } \mu \in \mathbf{P}(X) \text{ and } A \subset Y,$$

defines a Borel function $f_{\#} : \mathbf{P}(X) \rightarrow \mathbf{P}(Y)$. Moreover, if f is continuous, then so is $f_{\#}$.

(2) *The product of measures*

$$(\mu, \nu) \mapsto \mu \times \nu \quad \text{whenever } \mu \in \mathbf{P}(X) \text{ and } \nu \in \mathbf{P}(Y)$$

defines a continuous function $\mathbf{P}(X) \times \mathbf{P}(Y) \rightarrow \mathbf{P}(X \times Y)$.

Proof. We first prove (1). By [Kel75, Chapter 4, Theorem 16], [MS25, 2.11], and [Fed69, 2.2.2], $f_{\#}\mu$ is a Radon measure over Y . Note that it is enough to show $\mu \mapsto \int g d(f_{\#}\mu)$ is a Borel function whenever $g \in \mathcal{K}(Y)$ by [Men16b, 2.23]. Since $\int g d(f_{\#}\mu) = \int (g \circ f) d\mu$ whenever $\mu \in \mathbf{P}(X)$ and $g : Y \rightarrow \mathbf{R}$ is a bounded Borel function by [Fed69, 2.4.18], it reduces to show $\mu \mapsto \int h d\mu$ is a Borel function whenever $h : X \rightarrow \mathbf{R}$ is a bounded Borel function.

For $i \in \mathcal{P}$, let B_i be the family of all Borel functions $h : X \rightarrow \mathbf{R}$ with $\sup \text{im } |h| \leq i$ such that $\mu \mapsto \int h d\mu$ is a Borel function on $\mathbf{P}(X)$. By 3.8, the set B_i contains all continuous functions from X into $\mathbf{R} \cap \{r : |r| \leq i\}$. By [Fed69, 2.4.9] and employing the terminology of [Fed69, 2.2.15], B_i is a Baire class, hence contains all Baire functions with image contained in $\mathbf{R} \cap \{r : |r| \leq i\}$; applying [Fed69, 2.2.15] with $Y = \mathbf{R} \cap \{r : |r| \leq i\}$, we see B_i contains all Borel functions $h : X \rightarrow \mathbf{R}$ with $\sup \text{im } |g| \leq i$. Therefore, the main conclusion of (1) follows. For the postscript of (1), it is enough to show $\mu \mapsto \int g d(f_{\#}\mu) = \int (g \circ f) d\mu$ is continuous whenever $g \in \mathcal{K}(Y)$, which is immediate from 3.8.

Next, we will show (2). By Fubini's theorem [Fed69, 2.6.2], $\mu \times \nu$ is a Borel regular measure, hence by [Fed69, 2.2.2], a Radon measure whenever $\mu \in \mathbf{P}(X)$ and $\nu \in \mathbf{P}(Y)$. It is enough to show that $e_{\phi}(\mu, \nu) = \int \phi d(\mu \times \nu)$ defines a continuous function on $\mathbf{P}(X) \times \mathbf{P}(Y)$ whenever $\phi \in \mathcal{K}(X \times Y)$. If $\phi \in \mathcal{K}(X \times Y)$ satisfies $\phi(x, y) = \alpha(x)\beta(y)$ whenever $(x, y) \in X \times Y$ for some $\alpha \in \mathcal{K}(X)$ and $\beta \in \mathcal{K}(Y)$, then e_{ϕ} is continuous. Therefore, in view of 3.6, for each $\phi \in \mathcal{K}(X \times Y)$, e_{ϕ} is the uniform limit of a sequence of continuous functions, hence itself a continuous function. \square

Remark 4.20. The two statements in 4.19 are variants of [Kec95, 17.28, 17.40].

Definition 4.21. Suppose Y and Z are second-countable locally compact Hausdorff spaces, μ is a Radon measure over a locally compact Hausdorff space X , $f : Y \rightarrow Z$ is a Borel function, and g is a μ Young function of type Y . Then, we define the μ Young function $f_{\#}g$ of type Z by $(f_{\#}g)(x) = f_{\#}(g(x))$ whenever $x \in \text{dmn } g$.

Definition 4.22. Suppose X , Y , and Z are locally compact Hausdorff spaces, Y and Z are second-countable, μ is a Radon measure over X , and f and g are μ Young functions of types Y and Z , respectively. Then, we define the μ Young function $f \times g$ of type $Y \times Z$ by $(f \times g)(x) = f(x) \times g(x)$ whenever $x \in \text{dmn } f \cap \text{dmn } g$.

We finish this section by introducing the sum of two vector-valued functions and the product of two real-valued functions in the context of Young functions.

Definition 4.23 (see [Kle20, 14.20]). Suppose μ and ν are measures over a vector space Y , and $A : Y \times Y \rightarrow Y$ is defined by

$$A(x, y) = x + y \quad \text{whenever } x, y \in Y.$$

Then, the measure $\mu * \nu = A_{\#}(\mu \times \nu)$ is termed the *convolution of μ and ν* .

Remark 4.24 (see [Kle20, 14.21]). Suppose Y is a finite-dimensional Banach space. We readily verify the following basic properties of convolutions of measures.

- (1) Whenever $\mu, \nu \in \mathbf{P}(Y)$, we have $\mu * \nu = \nu * \mu$.
- (2) If $\mu, \nu \in \mathbf{P}(Y)$, then $\mu * \nu \in \mathbf{P}(Y)$.
- (3) If $\mu, \nu \in \mathbf{P}(Y)$ satisfy $\mu = \sum_{x \in A} f(x) \delta_x$ and $\nu = \sum_{y \in B} g(y) \delta_y$ for some finite subsets A and B of Y and functions $f : A \rightarrow \mathbf{R}$ and $g : B \rightarrow \mathbf{R}$, then $\mu * \nu = \sum_{(x, y) \in A \times B} f(x)g(y) \delta_{x+y}$.

Corollary 4.25. *Suppose Y is a finite-dimensional Banach space and d is as in 4.7. Then, the convolution of measures*

$$(\mu, \nu) \mapsto \mu * \nu \quad \text{whenever } \mu, \nu \in \mathbf{P}(Y)$$

defines a continuous function $\mathbf{P}(Y) \times \mathbf{P}(Y) \rightarrow \mathbf{P}(Y)$ with respect to the weak topologies. Furthermore, we have

$$d(\mu * \nu, \lambda * \eta) \leq d(\mu, \lambda) + d(\nu, \eta) \quad \text{whenever } \mu, \nu, \lambda, \eta \in \mathbf{P}(Y);$$

in particular, the convolution of measures defines a Lipschitzian map $(\mathbf{P}(Y) \times \mathbf{P}(Y), \rho) \rightarrow (\mathbf{P}(Y), d)$ where ρ satisfies $\rho((\mu, \nu), (\lambda, \eta)) = d(\mu, \nu) + d(\lambda, \eta)$ whenever $\mu, \nu, \lambda, \eta \in \mathbf{P}(Y)$.

Proof. The main assertion follows from 4.19(1)(2). The postscript follows from 4.9, 4.24(1), and the estimate

$$\begin{aligned} \int \gamma d(\mu * \lambda) - \int \gamma d(\nu * \lambda) &= \int (\int \gamma(x + y) d\mu x - \int \gamma(x + y) d\nu x) d\lambda y \\ &\leq d(\mu, \nu) \end{aligned}$$

whenever $\mu, \nu, \lambda \in \mathbf{P}(Y)$ and $\gamma \in \mathcal{K}(Y)$ with $\text{Lip } \gamma \leq 1$. □

Remark 4.26. Similarly, if $Y = \mathbf{R}$, replacing A with the multiplication on \mathbf{R} induces a continuous function $\mathbf{P}(\mathbf{R}) \times \mathbf{P}(\mathbf{R}) \rightarrow \mathbf{P}(\mathbf{R})$.

Definition 4.27. Suppose X is a locally compact Hausdorff space, Y is a finite-dimensional Banach space, μ is a Radon measure over X , and f and g are μ Young functions of type Y . Then, we define the μ Young function $f * g$ by $(f * g)(x) = f(x) * g(x)$ whenever $x \in \text{dmn } f \cap \text{dmn } g$.

Remark 4.28. By the postscript of 4.25, we have $\text{Lip } f * g \leq \text{Lip } f + \text{Lip } g$.

5 The test function spaces

Hypotheses. In this section, we always assume Y is a finite-dimensional Banach space with $\dim Y \geq 1$, $n \in \mathcal{P}$, and U is an open subset of \mathbf{R}^n .

The goal of this section is to prove an embedding theorem of $\mathcal{K}(X, \mathbf{E}(Y))$ into $\mathcal{K}(X \times Y, \text{Hom}(Y, \mathbf{R}))$, see 5.16, and the embedding theorems for other test function spaces of interest follow immediately. For this purpose, we first present basic results about the function space $\mathcal{K}(X, Z)$.

Lemma 5.1. *Suppose X is a locally compact Hausdorff space and W and Z are locally convex Hausdorff spaces. Then, the following four statements hold.*

- (1) *If $f : W \rightarrow Z$ is a continuous linear map, then post-composition with f defines a continuous linear map $\mathcal{K}(X, W) \rightarrow \mathcal{K}(X, Z)$.*
- (2) *If Z is the product of finitely many locally convex spaces Z_1, Z_2, \dots, Z_n , then we have the isomorphism of locally convex spaces*

$$\mathcal{K}(X, Z) \simeq \prod_{i=1}^n \mathcal{K}(X, Z_i);$$

in particular, we have $\mathcal{K}(X, Z^n) \simeq \mathcal{K}(X, Z)^n$.

- (3) *If Z is normed and K is a compact subset of X , then the topology of $\mathcal{K}_K(X, Z)$ is given by the norm with value $\sup \text{im } |f|$ at $f \in \mathcal{K}_K(X, Z)$.*
- (4) *If K_i is a sequence of compact subsets of X such that $K_i \subset \text{Int } K_{i+1}$ for $i \in \mathcal{P}$ and $X = \bigcup_{i=1}^{\infty} K_i$, then $\mathcal{K}(X, Z)$ is the strict inductive limit of $\mathcal{K}_{K_i}(X, Z)$ as $i \rightarrow \infty$.*

Proof. It is straightforward to verify (1) from the definitions. The statements (2) and (3) are proved in [Bou04a, III, §1, No. 1]. To prove (4), by 2.9, it is enough to check the inclusion map $\mathcal{K}_{K_i}(X, Z) \rightarrow \mathcal{K}_{K_j}(X, Z)$ is a homeomorphic embedding whenever $i \leq j$, and this is straightforward from the definitions. \square

Lemma 5.2. *Suppose X is a second-countable locally compact Hausdorff space, K is a compact subset of X , and Z is a separable Banach space. Then, the function space $\mathcal{K}_K(X, Z)$ is a separable Banach space.*

Proof. Since Z is a Banach space and $\mathcal{K}_K(X, Z)$ is endowed with the supremum norm, it is straightforward to check that $\mathcal{K}_K(X, Z)$ is a Banach space. Since $\mathcal{K}_K(X)$ is separable by [Men16b, 2.2, 2.23], it follows from 3.5 that $\mathcal{K}_K(X, Z)$ is separable. \square

Lemma 5.3. *Suppose Z is a normed space, X is a locally compact Hausdorff space, K_i is a sequence of compact subsets of X such that $K_0 = \emptyset$, $K_{i-1} \subset \text{Int } K_i$ whenever $i \in \mathcal{P}$, and $X = \bigcup_{i=1}^{\infty} K_i$. Then, the subsets*

$$V_\alpha = \mathcal{K}(X, Z) \cap \{f : |f(x)| \leq \alpha(i)^{-1} \text{ for } i \in \mathcal{P} \text{ and } x \in X \sim K_{i-1}\}$$

corresponding to each $\alpha \in \mathcal{P}^{\mathcal{P}}$ form a fundamental system of neighborhoods of 0 in $\mathcal{K}(X, Z)$.

Proof. Clearly, V_α is a neighborhood of 0 in $\mathcal{K}(X, Z)$ whenever $\alpha \in \mathcal{P}^{\mathcal{P}}$. Let V be a convex neighborhood of 0 in $\mathcal{K}(X, Z)$. There exists $\beta \in \mathcal{P}^{\mathcal{P}}$ such that

$$\mathbf{B}(0, \beta(i)^{-1}) \cap \mathcal{K}_{K_i}(X, Z) \subset V \cap \mathcal{K}_{K_i}(X, Z) \quad \text{whenever } i \in \mathcal{P}.$$

Choose non-negative functions $\phi_i \in \mathcal{K}(X)$ for $i \in \mathcal{P}$ such that

$$\text{spt } \phi_i \subset (\text{Int } K_{i+1}) \sim K_{i-1} \quad \text{and} \quad \sum_{j=1}^{\infty} 2^{-j} \phi_j(x) = 1$$

whenever $i \in \mathcal{P}$ and $x \in X$, and choose $\alpha \in \mathcal{P}^{\mathcal{P}}$ such that

$$\alpha(i)^{-1} \sup \text{im } |\phi_i| \leq \beta(i+1)^{-1} \quad \text{whenever } i \in \mathcal{P}.$$

Note that if $i, j \in \mathcal{P}$ and $f \in V_\alpha \cap \mathcal{K}_{K_i}(X, Z)$, then

$$\text{spt } \phi_j f \subset (\text{Int } K_{j+1}) \sim K_{j-1} \quad \text{and} \quad \sup \text{im } |\phi_j f| \leq \beta(j+1)^{-1};$$

therefore

$$\begin{aligned} \phi_j f &\in V \cap \mathcal{K}_{K_{j+1}}(X, Z), \quad \text{if } j \leq i, \\ \phi_j f &= 0, \quad \text{if } j > i. \end{aligned}$$

Since V is convex, $0 \in V$, and $f = \sum_{j=1}^i 2^{-j} (\phi_j f) + 2^{-i} \cdot 0$, we conclude that $f \in V$, hence $V_\alpha \subset V$. \square

Remark 5.4. The proof of 5.3 is adapted from the proof of [Sch66, p. 66, Théorème II].

Next, we will study the topological vector structure of $\mathbf{E}(Y)$.

Lemma 5.5. *The following three statements hold.*

(1) *The derivative defines a continuous monomorphism*

$$\mathbf{E}(Y) \rightarrow \mathcal{K}(Y, \text{Hom}(Y, \mathbf{R}))$$

and its restriction to $\mathbf{E}_s(Y)$ gives a norm-preserving embedding

$$\mathbf{E}_s(Y) \rightarrow \mathcal{K}_{\mathbf{B}(0,s)}(Y, \text{Hom}(Y, \mathbf{R}))$$

whenever $0 \leq s < \infty$.

(2) *$\mathbf{E}_s(Y)$ is a separable Banach space whenever $0 \leq s < \infty$.*

(3) *$\mathbf{E}(Y)$ is the strict inductive limit of $\mathbf{E}_i(Y)$ as $i \rightarrow \infty$.*

Proof. We first prove (1). From the definition, we see that the derivative defines a norm-preserving embedding $\mathbf{E}_s(Y) \rightarrow \mathcal{K}_{\mathbf{B}(0,s)}(Y, \text{Hom}(Y, \mathbf{R}))$ whenever $0 \leq s < \infty$, hence a continuous monomorphism $\mathbf{E}(Y) \rightarrow \mathcal{K}(Y, \text{Hom}(Y, \mathbf{R}))$ by the universal property 2.5. The second assertion then follows from 5.1(4) and 2.10(1).

Next, we aim to show (2). Let f_i be a Cauchy sequence in $\mathbf{E}_s(Y)$. Then, Df_i is a Cauchy sequence in $\mathcal{K}_{\mathbf{B}(0,s)}(Y, \text{Hom}(Y, \mathbf{R}))$; by 4.3, the sequence f_i is also Cauchy with respect to the supremum metric. Therefore, their pointwise limits $f : Y \rightarrow \mathbf{R}$ and $F : Y \rightarrow \text{Hom}(Y, \mathbf{R})$ are both continuous. By Taylor's theorem, we can show that Df exists and equals F . It follows that $\mathbf{E}_s(Y)$ is complete. The separability of $\mathbf{E}_s(Y)$ follows from (1) and 5.2.

Finally, since the inclusion map $\mathbf{E}_r(Y) \rightarrow \mathbf{E}_s(Y)$ is a norm-preserving embedding whenever $0 \leq r \leq s < \infty$, (3) follows from 2.9. \square

Remark 5.6. If $Y = \mathbf{R}$, then both maps in 5.5(1) are isomorphisms of locally convex spaces since the formula

$$\gamma(y) = \int_0^1 \langle y, f(ty) \rangle dt \quad \text{whenever } y \in \mathbf{R}$$

defines a member in $\mathbf{E}_s(\mathbf{R})$ whenever $f \in \mathcal{K}_{\mathbf{B}(0,s)}(\mathbf{R}, \text{Hom}(\mathbf{R}, \mathbf{R}))$ and $0 \leq s < \infty$. If $Y = \mathbf{R}^k$ with $k \geq 2$, then both maps in 5.5(1) are not surjective; in fact, we may choose a non-negative function $\omega \in \mathcal{D}(\mathbf{R}, \mathbf{R})$ such that $\omega(r) = 1$ for $|r| \leq 1$ and define

$$f(x) = \omega(|x|)(x_2, -x_1, 0, \dots, 0) \quad \text{whenever } x = (x_1, x_2, \dots, x_k) \in \mathbf{R}^k.$$

Then, there exists no $\gamma \in \mathbf{E}(\mathbf{R}^k)$ such that $D\gamma = f$; here we identify \mathbf{R}^k with $\text{Hom}(\mathbf{R}^k, \mathbf{R})$.

To show the map $\mathbf{E}(Y) \rightarrow \mathcal{K}(Y, \text{Hom}(Y, \mathbf{R}))$ as in 5.5(1) is still a homeomorphic embedding for $\dim Y \geq 2$, we need to study the fundamental system of neighborhoods of 0 in $\mathbf{E}(Y)$; for this purpose, we introduce another function space $\tilde{\mathbf{E}}(Y)$ that is isomorphic to $\mathbf{E}(Y)$ as locally convex spaces when $\dim Y \geq 2$, but its members have compact support.

Definition 5.7. We define

$$\tilde{\mathbf{E}}_s(Y) = \{\tilde{\gamma} : \tilde{\gamma} : Y \rightarrow \mathbf{R} \text{ is of class 1, } \text{spt } \tilde{\gamma} \subset \mathbf{B}(0, s)\} \quad \text{whenever } 0 \leq s < \infty$$

endowed with the norm $\tilde{\gamma} \mapsto \sup \text{im } \|D\tilde{\gamma}\|$, and the space

$$\tilde{\mathbf{E}}(Y) = \bigcup \{\tilde{\mathbf{E}}_s(Y) : 0 \leq s < \infty\}$$

is endowed with the locally convex final topology induced by the inclusion maps $\tilde{\mathbf{E}}_s(Y) \rightarrow \tilde{\mathbf{E}}(Y)$.

Remark 5.8. Similar arguments as in the proof of 5.5 show that $\tilde{\mathbf{E}}(Y)$ is the strict inductive limit of the separable Banach spaces $\tilde{\mathbf{E}}_i(Y)$ as $i \rightarrow \infty$.

Remark 5.9. The map $\tilde{\gamma} \rightarrow \tilde{\gamma} - \tilde{\gamma}(0)$ defines a norm-preserving embedding from $\tilde{\mathbf{E}}_s(Y)$ into $\mathbf{E}_s(Y)$ whenever $0 \leq s < \infty$, hence by 2.5 a continuous monomorphism from $\tilde{\mathbf{E}}(Y)$ into $\mathbf{E}(Y)$, and therefore by 5.1(1) post-composition defines a continuous monomorphism $I : \mathcal{K}(X, \tilde{\mathbf{E}}(Y)) \rightarrow \mathcal{K}(X, \mathbf{E}(Y))$ whenever X is a locally compact Hausdorff space. For $\dim Y \geq 2$, these maps are isomorphisms of locally convex spaces.

Lemma 5.10. *Suppose $L_0 = \emptyset$ and $L_i = \mathbf{B}(0, i) \cap Y$ for $i \in \mathcal{P}$. Then, the subsets \widetilde{W}_α consisting of $\tilde{\gamma} \in \widetilde{\mathbf{E}}(Y)$ satisfying*

$$|\tilde{\gamma}(y)| + \|\mathbf{D}\tilde{\gamma}(y)\| \leq \alpha(i)^{-1} \quad \text{whenever } i \in \mathcal{P} \text{ and } y \in Y \sim L_{i-1}$$

corresponding to each $\alpha \in \mathcal{P}^{\mathcal{P}}$ form a fundamental system of neighborhoods of 0 in $\widetilde{\mathbf{E}}(Y)$.

Proof. Clearly, the set \widetilde{W}_α is convex, symmetric, and absorbent whenever $\alpha \in \mathcal{P}^{\mathcal{P}}$. Note that if

$$\begin{aligned} i \in \mathcal{P}, \quad \varepsilon &= (2i)^{-1} \inf\{\alpha(j)^{-1} : j = 1, 2, \dots, i\} \\ \tilde{\gamma} &\in \mathbf{B}(0, \varepsilon) \cap \widetilde{\mathbf{E}}_i(Y), \quad v \in Y, \quad |v| = 1, \quad \text{and} \quad y(t) = (i-t)v, \end{aligned}$$

then we have

$$\begin{aligned} |\tilde{\gamma}(y(t))| &\leq \int_0^t | \langle -v, \mathbf{D}\tilde{\gamma}(y(s)) \rangle | \, d\mathcal{L}^1 s \\ &\leq \int_0^t \|\mathbf{D}\tilde{\gamma}(y(s))\| \, d\mathcal{L}^1 s \\ &\leq i\varepsilon \end{aligned}$$

whenever $0 \leq t \leq i$. Thus, $\mathbf{B}(0, \varepsilon) \cap \widetilde{\mathbf{E}}_i(Y) \subset \widetilde{W}_\alpha \cap \widetilde{\mathbf{E}}_i(Y)$, and we conclude \widetilde{W}_α is a neighborhood of 0 in $\widetilde{\mathbf{E}}(Y)$ whenever $\alpha \in \mathcal{P}^{\mathcal{P}}$.

Let V be a convex neighborhood of 0 in $\widetilde{\mathbf{E}}(Y)$. Then, there exists $\beta \in \mathcal{P}^{\mathcal{P}}$ such that

$$\mathbf{B}(0, \beta(i)^{-1}) \cap \widetilde{\mathbf{E}}_i(Y) \subset V \cap \widetilde{\mathbf{E}}_i(Y) \quad \text{whenever } i \in \mathcal{P}.$$

Choose non-negative functions $\phi_i : Y \rightarrow \mathbf{R}$ of class 1 for $i \in \mathcal{P}$ such that

$$\text{spt } \phi_i \subset (\text{Int } L_{i+1}) \sim L_{i-1} \quad \text{and} \quad \sum_{i=1}^{\infty} 2^{-i} \phi_i(y) = 1$$

whenever $i \in \mathcal{P}$ and $y \in Y$, and choose $\alpha \in \mathcal{P}^{\mathcal{P}}$ such that

$$\alpha(i)^{-1} \sup \text{im}(|\phi_i| + \|\mathbf{D}\phi_i\|) \leq \beta(i+1)^{-1} \quad \text{whenever } i \in \mathcal{P}.$$

Note that if $i, j \in \mathcal{P}$ and $\tilde{\gamma} \in \widetilde{W}_\alpha \cap \widetilde{\mathbf{E}}_i(Y)$, then

$$\text{spt } \phi_j \tilde{\gamma} \subset (\text{Int } L_{j+1}) \sim L_{j-1} \quad \text{and} \quad \sup \text{im} \|\mathbf{D}(\phi_j \tilde{\gamma})\| \leq \beta(j+1)^{-1};$$

therefore

$$\begin{aligned} \phi_j \tilde{\gamma} &\in V \cap \widetilde{\mathbf{E}}_{j+1}(Y), \quad \text{if } j \leq i, \\ \phi_j \tilde{\gamma} &= 0, \quad \text{if } j > i. \end{aligned}$$

Since V is convex, $0 \in V$, and $\tilde{\gamma} = \sum_{j=1}^i 2^{-j} (\phi_j \tilde{\gamma}) + 2^{-i} \cdot 0$, we conclude that $\tilde{\gamma} \in V$, hence that $\widetilde{W}_\alpha \subset V$. \square

Remark 5.11. Suppose X is a second-countable locally compact Hausdorff space. Whenever \widetilde{W}_α and V_α are as in 5.10 and 5.3, respectively, the images of $\widetilde{W}_\alpha \times \widetilde{W}_\alpha$ and $V_\alpha \times V_\alpha$ under multiplication are contained in \widetilde{W}_α and V_α , respectively. It follows from [Bou87, I, §1, No. 6, Proposition 5] that the maps $\widetilde{\mathbf{E}}(Y) \times \widetilde{\mathbf{E}}(Y) \rightarrow \widetilde{\mathbf{E}}(Y)$ and $\mathcal{K}(X) \times \mathcal{K}(X) \rightarrow \mathcal{K}(X)$ induced by multiplication are continuous.

To prove the embedding theorem, we still need several lemmas.

Lemma 5.12. *Suppose $\alpha \in \mathcal{P}^{\mathcal{P}}$. Then, there exists a positive convex decreasing function $h : \mathbf{R} \rightarrow \mathbf{R}$ of class 1 such that*

$$h(x) < \alpha(i)^{-1} \quad \text{whenever } x \geq i - 1 \text{ and } i \in \mathcal{P}.$$

Proof. Define β and δ in $\mathcal{P}^{\mathcal{P}}$ by

$$\begin{aligned} \beta(i) &= \sup\{\alpha(j) : 1 \leq j \leq i\} + i; \\ \delta(i) &= 2^{\beta(i)} \end{aligned}$$

whenever $i \in \mathcal{P}$. Note that δ satisfies

$$\alpha(i) < \delta(i) \leq \delta(i+1)/2 \quad \text{and} \quad \delta(i)^{-1} - \delta(i+1)^{-1} > \delta(i+1)^{-1} - \delta(i+2)^{-1}$$

whenever $i \in \mathcal{P}$. Then, the function $f : \mathbf{R} \rightarrow \mathbf{R}$ defined by

$$f(x) = \begin{cases} \frac{1}{\delta(1)} - \left(\frac{1}{\delta(1)} - \frac{1}{\delta(2)} \right) x & \text{if } x \leq 0 \\ \frac{1}{\delta(i)} - \left(\frac{1}{\delta(i)} - \frac{1}{\delta(i+1)} \right) (x - (i-1)) & \text{if } i \in \mathcal{P}, i-1 \leq x \leq i \end{cases}$$

is a convex decreasing function such that

$$f(x) < \alpha(i)^{-1} \quad \text{whenever } x \geq i - 1 \text{ and } i \in \mathcal{P}.$$

Choose a function $g : \mathbf{R} \rightarrow \mathbf{R}$ of class 1 such that $g \geq 0$, $\int g d\mathcal{L}^1 = 1$, and $\text{spt } g \subset \mathbf{R} \cap \{x : -1 \leq x \leq 0\}$.

Let h be the convolution $f * g$, and we will verify that h has the desired properties. Clearly, h is a positive function of class 1. Since

$$h(x) - f(x) = \int_{\mathbf{R} \cap \{y : -1 \leq y \leq 0\}} (f(x-y) - f(x))g(y) d\mathcal{L}^1 y < 0,$$

it follows that $h \leq f$. To show that h is decreasing, we compute

$$h(x+t) - h(x) = \int (f((x-y)+t) - f(x-y))g(y) d\mathcal{L}^1 y < 0$$

whenever $x \in \mathbf{R}$ and $t > 0$. For $x, z \in \mathbf{R}$ and $0 \leq \lambda \leq 1$, we compute

$$\begin{aligned} (f * g)(\lambda x + (1-\lambda)z) &= \int f(\lambda x + (1-\lambda)z - y)g(y) d\mathcal{L}^1 y \\ &= \int f(\lambda(x-y) + (1-\lambda)(z-y))g(y) d\mathcal{L}^1 y \\ &\leq \lambda \int f(x-y)g(y) d\mathcal{L}^1 y \\ &\quad + (1-\lambda) \int f(z-y)g(y) d\mathcal{L}^1 y \\ &= \lambda(f * g)(x) + (1-\lambda)(f * g)(z) \end{aligned}$$

and it follows that h is convex. \square

Lemma 5.13. *Suppose X is a locally compact Hausdorff space, K is a compact subset of X , and either $Y = \mathbf{R}$, $F_i = \mathcal{K}_{\mathbf{B}(0,i)}(\mathbf{R})$, and $F = \mathcal{K}(\mathbf{R})$, or $F_i = \tilde{\mathbf{E}}_i(Y)$ and $F = \tilde{\mathbf{E}}(Y)$. Then, the locally convex final topology \mathcal{T}_1 on $\mathcal{K}_K(X, F)$ induced by the inclusion maps $\mathcal{K}_K(X, F_i) \rightarrow \mathcal{K}_K(X, F)$ for $i \in \mathcal{P}$ is identical to the topology \mathcal{T}_2 of uniform convergence on $\mathcal{K}_K(X, F)$.*

Proof. Since the inclusion map

$$\mathcal{K}_K(X, F_i) \rightarrow (\mathcal{K}_K(X, F), \mathcal{T}_2)$$

is continuous whenever $i \in \mathcal{P}$ by 5.1(1), it follows that $\mathcal{T}_2 \subset \mathcal{T}_1$.

Conversely, let V be a convex neighborhood of 0 in $(\mathcal{K}_K(X, F), \mathcal{T}_1)$. Then, there exists $\beta \in \mathcal{P}^{\mathcal{P}}$ such that

$$\mathbf{B}(0, \beta(i)^{-1}) \cap \mathcal{K}_K(X, F_i) \subset V \cap \mathcal{K}_K(X, F_i) \quad \text{whenever } i \in \mathcal{P}.$$

Let $L_0 = \emptyset$ and $L_i = \mathbf{B}(0, i) \cap Y$ for $i \in \mathcal{P}$. Choose non-negative functions $\phi_i : Y \rightarrow \mathbf{R}$ of class 1 such that

$$\text{spt } \phi_i \subset (\text{Int } L_{i+1}) \sim L_{i-1} \quad \text{and} \quad \sum_{i=1}^{\infty} 2^{-i} \phi(y) = 1$$

whenever $i \in \mathcal{P}$ and $y \in Y$, and choose $\alpha \in \mathcal{P}^{\mathcal{P}}$ such that

$$\alpha(i)^{-1} \sup \text{im}(|\phi_i| + \|\mathbf{D} \phi_i\|) \leq \beta(i+1)^{-1} \quad \text{whenever } i \in \mathcal{P}.$$

Let $\eta \in \mathcal{K}_K(X, F)$ such that $\text{im } \eta \subset U_\alpha$, where U_α equals either V_α or \widetilde{W}_α as in 5.3 with X and Z both replaced by \mathbf{R} and 5.10, respectively. By 5.1(4), 5.8, and 2.10(2), there exists $i \in \mathcal{P}$ such that $\text{im } \eta \subset F_i$. Since multiplication by ϕ_j induces a continuous linear map from F into F_{j+1} by 5.11, 5.1(4), 5.8, and 2.10(1) the function η_j defined by $\eta_j(x) = \phi_j \eta(x)$ for $x \in X$ belongs to $\mathcal{K}_K(X, F_{j+1})$ by 5.1(1). Observe that

$$\sup \text{im} |\eta_j| \leq \beta(j+1)^{-1}$$

whenever $j \in \mathcal{P}$. Note that $\eta_j = 0$ for $j > i$. Therefore, we have

$$\eta = \sum_{j=1}^i 2^{-j} \eta_j + 2^{-1} \cdot 0 \in V$$

since V is convex and $0 \in V$. It follows that $\mathcal{T}_1 \subset \mathcal{T}_2$. □

Lemma 5.14. *Suppose X is a locally compact Hausdorff space, K_i is a sequence of compact subsets of X such that $X = \bigcup_{i=1}^{\infty} K_i$ and $K_i \subset \text{Int } K_{i+1}$ whenever $i \in \mathcal{P}$, and either $Y = \mathbf{R}$, $F_i = \mathcal{K}_{\mathbf{B}(0,i)}(\mathbf{R})$, and $F = \mathcal{K}(\mathbf{R})$, or $F_i = \widetilde{\mathbf{E}}_i(Y)$ and $F = \widetilde{\mathbf{E}}(Y)$.*

Then, the following three statements hold.

(1) *Whenever K is a compact subset of X and $j \in \mathcal{P}$, we have*

$$\begin{aligned} \mathcal{K}_K(X, F) &= \bigcup_{i=1}^{\infty} \mathcal{K}_K(X, F) \cap \{f : \text{im } f \subset F_i\}, \\ \mathcal{K}_K(X, F) \cap \{f : \text{im } f \subset F_j\} &= \mathcal{K}_K(X, F_j). \end{aligned}$$

Consequently, $\mathcal{K}_K(X, F)$ is the strict inductive limit of $\mathcal{K}_K(X, F_i)$ as $i \rightarrow \infty$.

(2) $\mathcal{H}(X, F)$ is the strict inductive limit of $\mathcal{H}_{K_i}(X, F_i)$ as $i \rightarrow \infty$. Moreover, this statement remains true with F and F_i replaced by $\mathbf{E}(Y)$ and $\mathbf{E}_i(Y)$, respectively.

(3) The map $\iota : \mathcal{H}(X, \mathbf{E}(Y)) \rightarrow \mathcal{H}(X \times Y, \text{Hom}(Y, \mathbf{R}))$ defined by

$$\iota(\eta)(x, y) = (\mathbf{D}\eta(x))(y) \quad \text{for } \eta \in \mathcal{H}(X, \mathbf{E}(Y)) \text{ and } (x, y) \in X \times Y$$

is a continuous monomorphism; moreover, its restriction gives a norm-preserving embedding

$$\mathcal{H}_K(X, \mathbf{E}_s(Y)) \rightarrow \mathcal{H}_{K \times \mathbf{B}(0, s)}(X \times Y, \text{Hom}(Y, \mathbf{R}))$$

whenever K is a compact subset of X and $0 \leq s < \infty$.

Proof. The main assertion of (1) follows from 5.1(4), 5.8, and 2.10(1)(2). Therefore, the postscript of (1) follows from 2.9 and 5.13.

Next, we will prove (2). By the postscript of (1) and 2.11, we see the locally convex final topology on $\mathcal{H}(X, F)$ induced by the inclusion maps

$$\mathcal{H}_{K_i}(X, F) \rightarrow \mathcal{H}(X, F) \quad \text{for } i \in \mathcal{P}$$

is identical to the locally convex final topology induced by the inclusion maps

$$\mathcal{H}_{K_i}(X, F_j) \rightarrow \mathcal{H}(X, F) \quad \text{for } i, j \in \mathcal{P}.$$

On the other hand, since the inclusion map $\mathcal{H}_{K_i}(X, F_j) \rightarrow \mathcal{H}_{K_k}(X, F_k)$ is a homeomorphic embedding whenever $i, j \in \mathcal{P}$ and $k = \sup\{i, j\}$, the main assertion of (2) follows from 2.9. To prove the postscript of (2), it is enough to note that for $\dim Y = 1$ we have $\mathcal{H}_{\mathbf{B}(0, i)}(\mathbf{R}) \simeq \mathbf{E}_i(\mathbf{R})$ and $\mathcal{H}(\mathbf{R}) \simeq \mathbf{E}(\mathbf{R})$ by 5.6, and for $\dim Y \geq 2$, we have $\tilde{\mathbf{E}}_s(Y) \simeq \mathbf{E}_s(Y)$ and $\tilde{\mathbf{E}}(Y) \simeq \mathbf{E}(Y)$ by 5.9.

Finally, we aim to prove (3). Whenever K is a compact subset of X and $0 \leq s < \infty$, by 5.5(1) and 5.1(1), post-composition with the derivative induces a norm-preserving embedding

$$\mathcal{H}_K(X, \mathbf{E}_s(Y)) \rightarrow \mathcal{H}_K(X, \mathcal{H}_{\mathbf{B}(0, s)}(Y, \text{Hom}(Y, \mathbf{R}))),$$

and it follows from 3.3 that so is

$$\mathcal{H}_K(X, \mathbf{E}_s(Y)) \rightarrow \mathcal{H}_{K \times \mathbf{B}(0, s)}(X \times Y, \text{Hom}(Y, \mathbf{R})).$$

These embeddings then induce a continuous linear map that agrees with ι by (2) and 2.5. The postscript follows from 5.1(4) and 2.10(1). \square

Remark 5.15. If $\dim Y = 1$, then the embeddings in 5.14(3) are norm-preserving isomorphisms and hence ι is an isomorphism of locally convex spaces. If $\dim Y \geq 2$, then the map $I : \mathcal{H}(X, \tilde{\mathbf{E}}(Y)) \rightarrow \mathcal{H}(X, \mathbf{E}(Y))$ as in 5.9 is an isomorphism; it follows that ι is a homeomorphic embedding if and only if so is the composition $\tilde{\iota} = \iota \circ I$.

The following theorem shows that the map $\tilde{\iota}$ is a homeomorphic embedding (even for the case $\dim Y = 1$); therefore, so is ι .

Theorem 5.16. *Suppose X is a locally compact Hausdorff space, $K_0 = \emptyset$, K_i is a sequence of compact subsets of X such that $K_i \subset \text{Int } K_{i+1}$ whenever $i \in \mathcal{P}$ and $\bigcup_{i=1}^{\infty} K_i = X$, let $L_0 = \emptyset$ and $L_i = \mathbf{B}(0, i) \cap Y$ whenever $i \in \mathcal{P}$, and let $C_i = K_i \times L_i$ whenever $i \in \mathcal{P} \cup \{0\}$.*

Then, the subsets \widetilde{W}_α consisting of $\eta \in \mathcal{K}(X, \widetilde{\mathbf{E}}(Y))$ satisfying

$$|\eta(x)(y)| + \|\mathbf{D}(\eta(x))(y)\| \leq \alpha(i)^{-1}$$

whenever $i \in \mathcal{P}$ and $(x, y) \in (X \times Y) \sim C_{i-1}$, corresponding to each $\alpha \in \mathcal{P}^{\mathcal{P}}$ form a fundamental system of neighborhoods of 0 of $\mathcal{K}(X, \widetilde{\mathbf{E}}(Y))$. Furthermore, the map $\tilde{\iota} : \mathcal{K}(X, \widetilde{\mathbf{E}}(Y)) \rightarrow \mathcal{K}(X \times Y, \text{Hom}(Y, \mathbf{R}))$ defined by

$$\tilde{\iota}(\eta)(x, y) = \mathbf{D}(\eta(x))(y) \quad \text{for } (x, y) \in X \times Y$$

is a homeomorphic embedding.

Proof. Clearly, the set \widetilde{W}_α is convex and symmetric; by 5.14(1), \widetilde{W}_α is absorbent whenever $\alpha \in \mathcal{P}^{\mathcal{P}}$. Note that if

$$\begin{aligned} i \in \mathcal{P}, \quad \varepsilon &= (2i)^{-1} \inf\{\alpha(j)^{-1} : j = 1, 2, \dots, i\}, \\ x \in X, \quad v \in Y, \quad |v| &= 1, \quad \text{and} \quad y(t) = (i-t)v, \\ \eta &\in \mathbf{B}(0, \varepsilon) \cap \mathcal{K}_{K_i}(X, \widetilde{\mathbf{E}}_i(Y)) \end{aligned}$$

then we have

$$\begin{aligned} |\eta(x)(y(t))| &\leq \int_0^t | \langle -v, \mathbf{D}(\eta(x))(y(s)) \rangle | \, d\mathcal{L}^1 s \\ &\leq \int_0^t \|\mathbf{D}(\eta(x))(y(s))\| \, d\mathcal{L}^1 s \\ &\leq t\varepsilon \end{aligned}$$

whenever $0 \leq t \leq i$. Thus, $\mathbf{B}(0, \varepsilon) \cap \mathcal{K}_{K_i}(X, \widetilde{\mathbf{E}}_i(Y)) \subset \widetilde{W}_\alpha \cap \mathcal{K}_{K_i}(X, \widetilde{\mathbf{E}}_i(Y))$, and we conclude \widetilde{W}_α is a neighborhood of 0 in $\mathcal{K}(X, \widetilde{\mathbf{E}}(Y))$ whenever $\alpha \in \mathcal{P}^{\mathcal{P}}$.

Let V be a convex neighborhood of 0 in $\mathcal{K}(X, \widetilde{\mathbf{E}}(Y))$. Then, by 5.14(2), there exists $\beta \in \mathcal{P}^{\mathcal{P}}$ such that

$$\mathbf{B}(0, \beta(i)^{-1}) \cap \mathcal{K}_{K_i}(X, \widetilde{\mathbf{E}}_i(Y)) \subset V \cap \mathcal{K}_{K_i}(X, \widetilde{\mathbf{E}}_i(Y)) \quad \text{whenever } i \in \mathcal{P}.$$

Choose non-negative continuous functions $f_i : X \rightarrow \mathbf{R}$ and non-negative functions $g_i : Y \rightarrow \mathbf{R}$ of class 1 for $i \in \mathcal{P}$ such that

$$\text{spt } f_i \subset (\text{Int } K_{i+1}) \sim K_{i-1} \quad \text{and} \quad \text{spt } g_i \subset (\text{Int } L_{i+1}) \sim L_{i-1}$$

whenever $i \in \mathcal{P}$ and such that

$$\sum_{i=1}^{\infty} f_i(x) = 1 \quad \text{and} \quad \sum_{i=1}^{\infty} g_i(y) = 1$$

whenever $(x, y) \in X \times Y$. For $i \in \mathcal{P}$, we define a continuous function ϕ_i with values in $\widetilde{\mathbf{E}}(Y)$ by

$$\phi_i(x) = 2^i \left(\left(\sum_{j=1}^i f_j(x) \right) \left(\sum_{j=1}^i g_j \right) - \left(\sum_{j=1}^{i-1} f_j(x) \right) \left(\sum_{j=1}^{i-1} g_j \right) \right) \quad \text{for } x \in X.$$

Note that

$$(X \times Y) \cap \{(x, y) : \phi_i(x)(y) \neq 0\} \subset (\text{Int } C_{i+1}) \sim C_{i-1} \quad \text{for } i \in \mathcal{P}$$

and that

$$\sum_{i=1}^{\infty} 2^{-i} \phi_i(x)(y) = 1 \quad \text{for } (x, y) \in X \times Y.$$

Choose $\alpha \in \mathcal{P}^{\mathcal{P}}$ such that

$$\alpha(i)^{-1} (|\phi_i(x)(y)| + \|D(\phi_i(x))(y)\|) \leq \beta(i+1)^{-1}$$

whenever $(x, y) \in X \times Y$ and $i \in \mathcal{P}$. If $i, j \in \mathcal{P}$ and $\eta \in \widetilde{W}_\alpha \cap \mathcal{K}_{K_i}(X, \widetilde{\mathbf{E}}_i(Y))$, then the function $\phi_j \eta$ defined by

$$(\phi_j \eta)(x) = \phi_j(x) \eta(x) \quad \text{whenever } x \in X$$

satisfies $\phi_j \eta \in \mathcal{K}_{K_{j+1}}(X, \widetilde{\mathbf{E}}_{j+1}(Y))$ because of 5.11 and 5.8, and

$$\sup\{\sup \text{im } \|D((\phi_j \eta)(x))\| : x \in X\} \leq \beta(j+1)^{-1};$$

therefore

$$\begin{aligned} \phi_j \eta &\in V \cap \mathcal{K}_{K_{j+1}}(X, \widetilde{\mathbf{E}}_{j+1}(Y)) \quad \text{if } j \leq i, \\ \phi_j \eta &= 0 \quad \text{if } j > i. \end{aligned}$$

Since V is convex, $0 \in V$, and $\eta = \sum_{j=1}^i 2^{-j} (\phi_j \eta) + 2^{-i} \cdot 0$, we conclude that $\eta \in V$, hence $\widetilde{W}_\alpha \subset V$.

Finally, by 5.15, we see \tilde{t} is a continuous monomorphism; hence, it remains to prove that \tilde{t}^{-1} is continuous. Let $\alpha \in \mathcal{P}^{\mathcal{P}}$ and let \widetilde{W}_α be a neighborhood of 0 in $\mathcal{K}(X, \widetilde{\mathbf{E}}(Y))$ defined as above. Now, we aim to find $\delta \in \mathcal{P}^{\mathcal{P}}$ such that

$$V_\delta \cap \text{im } \tilde{t} \subset \tilde{t}[\widetilde{W}_\alpha]$$

where V_δ is a neighborhood of 0 in $\mathcal{K}(X \times Y, \text{Hom}(Y, \mathbf{R}))$ as in 5.3 with X , K_i , and Z replaced by $X \times Y$, C_i , and $\text{Hom}(Y, \mathbf{R})$, respectively. By 5.12, there exists a positive convex decreasing function $h : \mathbf{R} \rightarrow \mathbf{R}$ of class 1 such that $h(i-1) \leq 2^{-1} \alpha(i)^{-1}$ whenever $i \in \mathcal{P}$. Then, we define $\delta \in \mathcal{P}^{\mathcal{P}}$ to satisfy

$$\delta(i)^{-1} \leq \inf\{-h'(i), 2^{-1} \alpha(i)^{-1}\} \quad \text{whenever } i \in \mathcal{P}$$

and let V_δ be as in 5.3. If

$$\begin{aligned} i &\in \mathcal{P}, \quad \eta \in \tilde{t}^{-1}[V_\delta] \cap \mathcal{K}_{K_i}(X, \widetilde{\mathbf{E}}_i(Y)), \\ v &\in Y, \quad |v| = 1, \quad \text{and} \quad y(t) = (i-t)v, \end{aligned}$$

then we have

$$\begin{aligned} |\eta(x)(y(t))| &\leq \int_0^t | \langle -v, D(\eta(x))(y(s)) \rangle | d\mathcal{L}^1 s \\ &\leq \int_0^t \|D(\eta(x))(y(s))\| d\mathcal{L}^1 s \\ &\leq \int_0^t -h'(i-s) d\mathcal{L}^1 s \\ &\leq h(|y(t)|) \end{aligned}$$

whenever $x \in X$ and $0 \leq t \leq i$. It follows that $|\eta(x)(y)| \leq h(|y|)$ whenever $(x, y) \in X \times Y$, hence $\tilde{t}^{-1}[V_\delta] \cap \mathcal{K}_{K_i}(X, \widetilde{\mathbf{E}}_i(Y)) \subset \widetilde{W}_\alpha$. \square

Corollary 5.17. *Suppose X is a locally compact Hausdorff space, $K_0 = \emptyset$, K_i is a sequence of compact subsets of X such that $K_i \subset \text{Int } K_{i+1}$ whenever $i \in \mathcal{P}$ and $\bigcup_{i=1}^{\infty} K_i = X$, and let $C_i = K_i \times (\mathbf{B}(0, i) \cap Y)$ whenever $i \in \mathcal{P} \cup \{0\}$.*

Then, the map $\iota : \mathcal{K}(X, \mathbf{E}(Y)) \rightarrow \mathcal{K}(X \times Y, \text{Hom}(Y, \mathbf{R}))$ defined by

$$\iota(\eta)(x, y) = \text{D}(\eta(x))(y) \quad \text{for } (x, y) \in X \times Y$$

is a homeomorphic embedding. Furthermore, the subsets W_α consisting of $\eta \in \mathcal{K}(X, \mathbf{E}(Y))$ satisfying

$$\|\text{D}(\eta(x))(y)\| \leq \alpha(i)^{-1} \quad \text{whenever } i \in \mathcal{P} \text{ and } (x, y) \in (X \times Y) \sim C_{i-1}$$

corresponding to each $\alpha \in \mathcal{P}^{\mathcal{P}}$ form a fundamental system of neighborhoods of 0 of $\mathcal{K}(X, \mathbf{E}(Y))$.

Proof. It follows from 5.15, 5.16, and 5.3. □

Corollary 5.18. *The map $\mathbf{E}(Y) \rightarrow \mathcal{K}(Y, \text{Hom}(Y, \mathbf{R}))$ defined as in 5.5(1) is a homeomorphic embedding.*

Furthermore, the subsets W_α consisting of $\gamma \in \mathbf{E}(Y)$ satisfying

$$\|\text{D} \gamma(y)\| \leq \alpha(i)^{-1} \quad \text{whenever } i \in \mathcal{P} \text{ and } |y| > i - 1$$

corresponding to each $\alpha \in \mathcal{P}^{\mathcal{P}}$ form a fundamental system of neighborhoods of 0 of $\mathbf{E}(Y)$.

Proof. If $\dim Y = 1$, it is proved in 5.6, and the postscript follows from 5.3. If $\dim Y \geq 2$, it follows from 5.17 applied with X and $K_i = X$ for $i \in \mathcal{P}$ being a singleton. □

Corollary 5.19. *The dual map $\mathcal{K}(Y, \text{Hom}(Y, \mathbf{R}))^* \rightarrow \mathbf{E}(Y)^*$ is an epimorphism.*

Proof. It follows from 5.18 and the Hahn-Banach theorem [Bou87, II, §4, No. 1, Proposition 2]. □

Remark 5.20. The following example shows that the dual map in 5.19 is not injective for $\dim Y = k > 1$. Assume $Y = \mathbf{R}^k$. Whenever $\omega \in \mathcal{D}(\mathbf{R}, \mathbf{R})$, $0 \notin \text{spt } \omega'$, and L is a non-zero anti-symmetric endomorphism on \mathbf{R}^k , the functional $\mu \in \mathcal{K}(\mathbf{R}^k, \text{Hom}(\mathbf{R}^k, \mathbf{R}))^*$ defined by

$$\mu(\delta) = \int \langle X(y), \delta(y) \rangle \text{d}\mathcal{L}^k y \quad \text{for } \delta \in \mathcal{K}(\mathbf{R}^k, \text{Hom}(\mathbf{R}^k, \mathbf{R}))$$

belongs to the kernel of the dual map $\mathcal{K}(\mathbf{R}^k, \text{Hom}(\mathbf{R}^k, \mathbf{R}))^* \rightarrow \mathbf{E}(\mathbf{R}^k)^*$, where $X \in \mathcal{D}(\mathbf{R}^k, \mathbf{R}^k)$ is defined by

$$X(y) = \omega(|y|)L(y) \quad \text{for } y \in \mathbf{R}^k,$$

because $\text{div } X = 0$.

To unify the test function spaces from domain and codomain, we also introduce the test function space on the product.

Definition 5.21. Whenever C is a compact subset of $U \times Y$, the function space $\mathbf{H}_C(U \times Y, \mathbf{R}^n)$ consists of all functions $\eta : U \times Y \rightarrow \mathbf{R}^n$ such that

(1) $\eta(x, \cdot) \bullet v \in \mathbf{E}(Y)$ whenever $x \in U$ and $v \in \mathbf{R}^n$.

(2) The function

$$(x, y) \mapsto D(\eta(x, \cdot))(y) \in \text{Hom}(Y, \mathbf{R}^n)$$

is continuous with compact support in C .

We endow $\mathbf{H}_C(U \times Y, \mathbf{R}^n)$ with the norm (see 5.22) whose value equal

$$\sup\{\|D(\eta(x, \cdot))(y)\| : (x, y) \in U \times Y\} \quad \text{at } \eta \in \mathbf{H}_C(U \times Y, \mathbf{R}^n),$$

and endow

$$\mathbf{H}(U \times Y, \mathbf{R}^n) = \bigcup\{\mathbf{H}_C(U \times Y, \mathbf{R}^n) : C \text{ is a compact subset of } U \times Y\}$$

with the locally convex final topology induced by the inclusion maps

$$\mathbf{H}_C(U \times Y, \mathbf{R}^n) \rightarrow \mathbf{H}(U \times Y, \mathbf{R}^n).$$

Remark 5.22. We shall verify that $\mathbf{H}_C(U \times Y, \mathbf{R}^n)$ is a normed space. Let $\eta \in \mathbf{H}_C(U \times Y, \mathbf{R}^n)$. Note that 5.21(1) implies that $\eta(x, 0) = 0$ whenever $x \in U$. If $(x, y) \in U \times Y$, then we have

$$\eta(x, y) = \int_0^1 \langle y, D(\eta(x, \cdot))(ty) \rangle d\mathcal{L}^1 t;$$

in particular, η is continuous. If η satisfies

$$\sup\{\|D(\eta(x, \cdot))(y)\| : x \in U \text{ and } y \in Y\} = 0,$$

then $\eta = 0$. Thus, $\mathbf{H}_C(U \times Y, \mathbf{R}^n)$ is a normed space.

Lemma 5.23. *The following two statements hold.*

(1) *Whenever K is a compact subset of U and $0 \leq s < \infty$, we have the following isomorphisms of Banach spaces*

$$\mathbf{H}_{K \times \mathbf{B}(0, s)}(U \times Y, \mathbf{R}^n) \simeq \mathcal{K}_K(U, \mathbf{E}_s(Y))^n;$$

in particular, $\mathbf{H}_{K \times \mathbf{B}(0, s)}(U \times Y, \mathbf{R}^n)$ is a separable Banach space.

(2) *Whenever C_i is a sequence of compact subsets of $U \times Y$ such that $U \times Y = \bigcup_{i=1}^{\infty} C_i$ and $C_i \subset \text{Int } C_{i+1}$ for $i \in \mathcal{P}$, $\mathbf{H}(U \times Y, \mathbf{R}^n)$ is the strict inductive limit of $\mathbf{H}_{C_i}(U \times Y, \mathbf{R}^n)$ as $i \rightarrow \infty$.*

Proof. To prove (1), let K be a compact subset of U and $0 \leq s < \infty$. From 5.21(2), the map

$$\mathbf{H}_{K \times \mathbf{B}(0, s)}(U \times Y, \mathbf{R}^n) \rightarrow \mathcal{K}_{K \times \mathbf{B}(0, s)}(U \times Y, \text{Hom}(Y, \mathbf{R}^n))$$

defined by

$$\eta \mapsto [(x, y) \mapsto D(\eta(x, \cdot))(y)]$$

is a homeomorphic embedding; noting from 5.1(2) that

$$\mathcal{K}_{K \times \mathbf{B}(0, s)}(U \times Y, \text{Hom}(Y, \mathbf{R}^n)) \simeq \mathcal{K}_{K \times \mathbf{B}(0, s)}(U \times Y, \text{Hom}(Y, \mathbf{R}))^n,$$

so is the map

$$f : \mathbf{H}_{K \times \mathbf{B}(0,s)}(U \times Y, \mathbf{R}^n) \rightarrow \mathcal{K}_{K \times \mathbf{B}(0,s)}(U \times Y, \text{Hom}(Y, \mathbf{R}))^n.$$

On the other hand, recall from 5.14(3) that the map

$$J : \mathcal{K}_K(U, \mathbf{E}_s(Y)) \rightarrow \mathcal{K}_{K \times \mathbf{B}(0,s)}(U \times Y, \text{Hom}(Y, \mathbf{R}))$$

defined by

$$J(\eta)(x, y) = \text{D}(\eta(x))(y) \quad \text{whenever } (x, y) \in U \times Y$$

is a homeomorphic embedding, and so is its Cartesian product $g = \prod_{i=1}^n J$. Since $\text{im } f = \text{im } g$, it follows that $g^{-1} \circ f$ defines an isomorphism between the normed spaces $\mathbf{H}_{K \times \mathbf{B}(0,s)}(U \times Y, \mathbf{R}^n)$ and $\mathcal{K}_K(U, \mathbf{E}_s(Y))^n$; in particular, $\mathbf{H}_{K \times \mathbf{B}(0,s)}(U \times Y, \mathbf{R}^n)$ is a separable Banach space by 5.5(2) and 5.2.

Since the inclusion map $\mathbf{H}_{C_i}(U \times Y, \mathbf{R}^n) \rightarrow \mathbf{H}_{C_j}(U \times Y, \mathbf{R}^n)$ is a norm preserving embedding whenever $i \leq j \in \mathcal{P}$, (2) follows from 2.9. \square

Remark 5.24. Suppose K_i is a sequence of compact subsets of U such that $U = \bigcup_{i=1}^{\infty} K_i$ and $K_i \subset \text{Int } K_{i+1}$ for $i \in \mathcal{P}$. By 5.1(1)(2), 2.17, and 5.14(2), we see $\mathcal{K}(U, \mathbf{E}(Y))^n \simeq \mathcal{K}(U, \mathbf{E}(Y)^n) \simeq \mathcal{K}(U, \mathbf{R}^n \otimes \mathbf{E}(Y))$ is the strict inductive limit of $\mathcal{K}_{K_i}(U, \mathbf{E}_i(Y))^n \simeq \mathcal{K}_{K_i}(U, \mathbf{E}_i(Y)^n) \simeq \mathcal{K}_{K_i}(U, \mathbf{R}^n \otimes \mathbf{E}_i(Y))$ as $i \rightarrow \infty$. Moreover, we have

$$\mathbf{H}(U \times Y, \mathbf{R}^n) \simeq \mathcal{K}(U, \mathbf{E}(Y))^n \simeq \mathcal{K}(U, \mathbf{E}(Y)^n) \simeq \mathcal{K}(U, \mathbf{R}^n \otimes \mathbf{E}(Y))$$

by 5.23(1)(2).

Theorem 5.25. *The map $\iota : \mathbf{H}(U \times Y, \mathbf{R}^n) \rightarrow \mathcal{K}(U \times Y, \text{Hom}(Y, \mathbf{R}^n))$ defined by*

$$\iota(\eta) = (\text{D} \eta(x, \cdot))(y) \quad \text{whenever } \eta \in \mathbf{H}(U \times Y, \mathbf{R}^n) \text{ and } (x, y) \in U \times Y$$

is a homeomorphic embedding.

Furthermore, if $K_0 = \emptyset$ and K_i is a sequence of compact subsets of U such that $U = \bigcup_{i=1}^{\infty} K_i$ and $K_i \subset \text{Int } K_{i+1}$ whenever $i \in \mathcal{P}$, then the subsets W_α consisting of $\eta \in \mathbf{H}(U \times Y, \mathbf{R}^n)$ satisfying

$$\sup \text{im } \|(\text{D} \eta(x, \cdot))(y)\| \leq \alpha(i)^{-1}$$

whenever $i \in \mathcal{P}$ and $(x, y) \in (U \times Y) \sim (K_{i-1} \times \mathbf{B}(0, i-1))$, corresponding to each $\alpha \in \mathcal{P}^{\mathcal{P}}$ form a fundamental system of neighborhoods of 0 of $\mathbf{H}(U \times Y, \mathbf{R}^n)$.

Proof. By 5.24, we have $\mathbf{H}(U \times Y, \mathbf{R}^n) \simeq \mathcal{K}(U, \mathbf{E}(Y))^n$. Since, by 5.17, 2.17, and 5.1(2), the composition

$$\mathcal{K}(U, \mathbf{E}(Y))^n \rightarrow \mathcal{K}(U \times Y, \text{Hom}(Y, \mathbf{R}))^n \simeq \mathcal{K}(U \times Y, \text{Hom}(Y, \mathbf{R}^n))$$

is a homeomorphic embedding and the main assertion follows. The postscript follows from 5.3. \square

Remark 5.26. In case $Y = \mathbf{R}$, the homeomorphic embedding

$$\mathbf{H}(U \times \mathbf{R}, \mathbf{R}^n) \rightarrow \mathcal{K}(U \times \mathbf{R}, \text{Hom}(\mathbf{R}, \mathbf{R}^n)) \simeq \mathcal{K}(U \times \mathbf{R}, \mathbf{R}^n)$$

is surjective; in fact, for $\phi \in \mathcal{K}(U \times \mathbf{R}, \mathbf{R}^n)$, the function $\eta : U \times \mathbf{R} \rightarrow \mathbf{R}^n$ defined by

$$\eta(x, y) = \int_0^y \phi(x, z) \, d\mathcal{L}^1 z$$

is a member in $\mathbf{H}(U \times \mathbf{R}, \mathbf{R}^n)$ whose image under the embedding equals ϕ .

Lemma 5.27. *Suppose A is a directed set, $(F_\alpha, f_{\beta\alpha})$ is an inductive system of locally convex spaces relative to A , $(D_\alpha, \delta_{\beta\alpha})$ is an inductive system of vector spaces relative to A , we denote by (F, f_α) and (D, δ_α) the inductive limit of $(F_\alpha, f_{\beta\alpha})$ and $(D_\alpha, \delta_{\beta\alpha})$ respectively, for $\alpha \in A$, D_α is a dense subspace of F_α and the inclusion map $\iota_\alpha : D_\alpha \rightarrow F_\alpha$ satisfies*

$$f_{\beta\alpha} \circ \iota_\alpha = \iota_\beta \circ \delta_{\beta\alpha} \quad \text{whenever } \alpha \leq \beta \in A.$$

Then, the vector space $D = \bigcup_{\alpha \in A} \delta_\alpha[D_\alpha]$ can be identified as a dense subspace of F . Moreover, if D_α is endowed with the subspace topology induced by ι_α , G is a complete Hausdorff locally convex space, and the family of continuous linear maps $g_\alpha : D_\alpha \rightarrow G$ for $\alpha \in A$ satisfies

$$g_\alpha = g_\beta \circ \delta_{\beta\alpha} \quad \text{whenever } \alpha \leq \beta \in A,$$

then the inductive limit of the unique extensions of g_α equals the unique extension of the inductive limit of g_α ; here, by convention, we also view G as the limit of the constant inductive system $(G, \mathbf{1}_G)$ of locally convex spaces relative to A .

Proof. From 2.7, the inductive limit ι of the inclusion maps $\iota_\alpha : D_\alpha \rightarrow F_\alpha$ is a monomorphism, and we identify D as its image under ι in F . Since D_α is dense in F_α , we have

$$\text{im } f_\alpha \subset \text{Clos im}(f_\alpha \circ \iota_\alpha) = \text{Clos im}(\iota \circ \delta_\alpha) \quad \text{whenever } \alpha \in A.$$

It follows that

$$\begin{aligned} F &= \bigcup \{\text{im } f_\alpha : \alpha \in A\} \subset \bigcup \{\text{Clos im}(\iota \circ \delta_\alpha) : \alpha \in A\} \\ &\subset \text{Clos} \bigcup \{\text{im}(\iota \circ \delta_\alpha) : \alpha \in A\} \\ &= \text{Clos } \iota[D]. \end{aligned}$$

To prove the postscript, we denote by h_α the unique continuous extension of g_α for $\alpha \in A$, and these h_α satisfy

$$h_\alpha = h_\beta \circ f_{\beta\alpha} \quad \text{whenever } \alpha \leq \beta \in A.$$

Then, the inductive limit h of h_α exists and satisfies

$$h \circ (\iota \circ \delta_\alpha) = h \circ (f_\alpha \circ \iota_\alpha) = g_\alpha \quad \text{whenever } \alpha \in A.$$

It follows that $h \circ \iota$ equals the inductive limit of g_α . Finally, if D is endowed with the subspace topology, then h is the unique continuous extension of the inductive limit of g_α . \square

Theorem 5.28. *Suppose $\mu : \mathcal{D}(U, \mathbf{R}^n) \otimes (\mathcal{E}(Y, \mathbf{R}) \cap \mathbf{E}(Y)) \rightarrow \mathbf{H}(U \times Y, \mathbf{R}^n)$ is the canonical map defined by*

$$\mu(\theta \otimes \gamma)(x, y) = \gamma(y)\theta(x)$$

for $\theta \in \mathcal{D}(U, \mathbf{R}^n)$, $\gamma \in \mathcal{E}(Y, \mathbf{R}) \cap \mathbf{E}(Y)$, and $(x, y) \in U \times Y$. Then, its image $\text{im } \mu$ is dense in $\mathbf{H}(U \times Y, \mathbf{R}^n)$.

Proof. Let K be a compact subset of U and $0 \leq s < \infty$. By 3.5, the image of the canonical monomorphism

$$\mathcal{K}_K(U) \otimes \mathbf{E}_s(Y) \rightarrow \mathcal{K}_K(U, \mathbf{E}_s(Y))$$

is dense; therefore, the image of its Cartesian product

$$(\mathcal{K}_K(U) \otimes \mathbf{E}_s(Y))^n \rightarrow \mathcal{K}_K(U, \mathbf{E}_s(Y))^n$$

is also dense. On the other hand, observe that

$$\mathcal{K}_K(U, \mathbf{R}^n) \otimes \mathbf{E}_s(Y) \simeq (\mathcal{K}_K(U) \otimes \mathbf{E}_s(Y))^n,$$

and by 5.23(1) we have $\mathcal{K}_K(U, \mathbf{E}_s(Y))^n \simeq \mathbf{H}_{K \times \mathbf{B}(0, s)}(U \times Y, \mathbf{R}^n)$ as Banach spaces. Thus, the composition

$$\mathcal{K}_K(U, \mathbf{R}^n) \otimes \mathbf{E}_s(Y) \rightarrow \mathbf{H}_{K \times \mathbf{B}(0, s)}(U \times Y, \mathbf{R}^n)$$

of the previous maps is a monomorphism whose image is dense. By convolution, we see $\mathcal{D}_K(U, \mathbf{R}^n)$ and $\mathcal{E}(Y, \mathbf{R}) \cap \mathbf{E}_s(Y)$ are dense in $\mathcal{K}_K(U, \mathbf{R}^n)$ and $\mathbf{E}_s(Y)$, respectively. Therefore, we may identify $\mathcal{D}_K(U, \mathbf{R}^n) \otimes (\mathcal{E}(Y, \mathbf{R}) \cap \mathbf{E}_s(Y))$ as a dense subspace of $\mathbf{H}_{K \times \mathbf{B}(0, s)}(U \times Y, \mathbf{R}^n)$.

Let K_i be a sequence of compact subset of U such that $U = \bigcup_{i=1}^{\infty} K_i$ and $K_i \subset \text{Int } K_{i+1}$ whenever $i \in \mathcal{P}$. Applying 5.27 with

$$A = \mathcal{P}, \quad D_i = \mathcal{D}_{K_i}(U, \mathbf{R}^n) \otimes (\mathcal{E}(Y, \mathbf{R}) \cap \mathbf{E}_i(Y)), \quad F_i = \mathbf{H}_{K_i \times \mathbf{B}(0, i)}(U \times Y, \mathbf{R}^n),$$

the assertion follows. \square

References

- [AFP00] Luigi Ambrosio, Nicola Fusco, and Diego Pallara. *Functions of bounded variation and free discontinuity problems*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 2000.
- [All72] William K. Allard. On the first variation of a varifold. *Ann. of Math. (2)*, 95:417–491, 1972.
- [Alm00] Frederick J. Almgren, Jr. *Almgren’s big regularity paper*, volume 1 of *World Scientific Monograph Series in Mathematics*. World Scientific Publishing Co., Inc., River Edge, NJ, 2000. Q -valued functions minimizing Dirichlet’s integral and the regularity of area-minimizing rectifiable currents up to codimension 2, With a preface by Jean E. Taylor and Vladimir Scheffer.

- [Bou87] Nicolas Bourbaki. *Topological vector spaces. Chapters 1–5*. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 1987. Translated from the French by H. G. Eggleston and S. Madan.
- [Bou89a] Nicolas Bourbaki. *Algebra. I. Chapters 1–3*. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 1989. Translated from the French, Reprint of the 1974 edition.
- [Bou89b] Nicolas Bourbaki. *General topology. Chapters 5–10*. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 1989. Translated from the French, Reprint of the 1966 edition.
- [Bou04a] Nicolas Bourbaki. *Integration. I. Chapters 1–6*. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2004. Translated from the 1959, 1965 and 1967 French originals by Sterling K. Berberian.
- [Bou04b] Nicolas Bourbaki. *Theory of sets*. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2004. Reprint of the 1968 English translation [Hermann, Paris; MR0237342].
- [DLS11] Camillo De Lellis and Emanuele Nunzio Spadaro. Q -valued functions revisited. *Mem. Amer. Math. Soc.*, 211(991):vi+79, 2011.
- [Fed69] Herbert Federer. *Geometric measure theory*. Die Grundlehren der mathematischen Wissenschaften, Band 153. Springer-Verlag New York, Inc., New York, 1969.
- [Fle57] W. H. Fleming. Functions with generalized gradient and generalized surfaces. *Ann. Mat. Pura Appl. (4)*, 44:92, 93–103, 1957.
- [Fle20] Wendell H. Fleming. Early developments in geometric measure theory. *Indiana Univ. Math. J.*, 69(1):5–36, 2020.
- [Hut86] John E. Hutchinson. Second fundamental form for varifolds and the existence of surfaces minimising curvature. *Indiana Univ. Math. J.*, 35(1):45–71, 1986.
- [Kec95] Alexander S. Kechris. *Classical descriptive set theory*, volume 156 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [Kel75] John L. Kelley. *General topology*. Graduate Texts in Mathematics, No. 27. Springer-Verlag, New York-Berlin, 1975. Reprint of the 1955 edition [Van Nostrand, Toronto, Ont.].
- [Kle20] Achim Klenke. *Probability theory—a comprehensive course*. Universitext. Springer, Cham, third edition, [2020] ©2020.
- [Men16a] Ulrich Menne. Sobolev functions on varifolds. *Proc. Lond. Math. Soc. (3)*, 113(6):725–774, 2016.
- [Men16b] Ulrich Menne. Weakly differentiable functions on varifolds. *Indiana Univ. Math. J.*, 65(3):977–1088, 2016.
- [MS25] Ulrich Menne and Christian Scharrer. A priori bounds for geodesic diameter. Part I. Integral chains with coefficients in a complete normed commutative group. *Rev. Mat. Iberoam.*, 41(1):29–72, 2025.

- [Sch66] Laurent Schwartz. *Théorie des distributions*. Publications de l'Institut de Mathématique de l'Université de Strasbourg, IX-X. Hermann, Paris, 1966. Nouvelle édition, entièrement corrigée, refondue et augmentée.
- [SW99] H. H. Schaefer and M. P. Wolff. *Topological vector spaces*, volume 3 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1999.
- [tD08] Tammo tom Dieck. *Algebraic topology*. EMS Textbooks in Mathematics. European Mathematical Society (EMS), Zürich, 2008.
- [Vil09] Cédric Villani. *Optimal transport*, volume 338 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 2009. Old and new.

AFFILIATION

Department of Mathematics
National Taiwan Normal University
No.88, Sec.4, Tingzhou Rd.
Wenshan Dist., TAIPEI CITY 116059
TAIWAN (R.O.C.)

EMAIL ADDRESS

hsinchuangchou@gmail.com